

Spectral estimates for matrix-valued periodic Dirac operators

Evgeny Korotyaev *

February 2, 2008

Abstract

We consider the first order periodic systems perturbed by a $2N \times 2N$ matrix-valued periodic potential on the real line. The spectrum of this operator is absolutely continuous and consists of intervals separated by gaps. We define the Lyapunov function, which is analytic on an associated N -sheeted Riemann surface. On each sheet the Lyapunov function has the standard properties of the Lyapunov function for the scalar case. The Lyapunov function has branch points, which we call resonances. We prove the existence of real or complex resonances. We determine the asymptotics of the periodic, anti-periodic spectrum and of the resonances at high energy (in terms of the Fourier coefficients of the potential). We show that there exist two types of gaps: i) stable gaps, i.e., the endpoints are periodic and anti-periodic eigenvalues, ii) unstable (resonance) gaps, i.e., the endpoints are resonances (real branch points). Moreover, we determine various new trace formulae for potentials and the Lyapunov exponent.

Keywords: periodic systems, spectrum, high energy asymptotics.

1 Introduction and main results

Consider the self-adjoint operator $\mathcal{K}$ acting on the space $L^2(\mathbb{R})^{2N}$ and given by

$$\mathcal{K}y = -iJ_1y' + V_t y, \quad J_1 = \begin{pmatrix} I_N & 0 \\ 0 & -I_N \end{pmatrix}, \quad V = \begin{pmatrix} 0 & v \\ v^* & 0 \end{pmatrix}, \quad N \geq 1,$$

here and below we use the notation $(') = \partial/\partial t$ and I_N is the identity $N \times N$ matrix; v is the **complex** 1-periodic $N \times N$ matrix and $V = V_t$ belongs to the Hilbert space $\mathcal{H}$ given by

$$\mathcal{H} = \left\{ V = \begin{pmatrix} 0 & v \\ v^* & 0 \end{pmatrix}, v = v^\top = \{v_{jk}(t)\}_{j,k=1}^N, \quad t \in \mathbb{R}/\mathbb{Z}, \quad \|V\|^2 = \int_0^1 \text{Tr } V_t^2 dt < \infty \right\}.$$

*Institut für Mathematik, Humboldt Universität zu Berlin, Rudower Chaussee 25, 12489, Berlin, Germany, e-mail: evgeny@math.hu-berlin.de

Note that $\operatorname{Re} v$ and $\operatorname{Im} v$ are self-adjoint. **Without loss of generality we assume**

$$\mathcal{V} = \int_0^1 V_t^2 dt = \mathcal{V}_0 \oplus \mathcal{V}_0, \quad \mathcal{V}_0 = \operatorname{diag}\{\nu_1, \dots, \nu_N\}, \quad 0 \leq \nu_1 \leq \nu_2 \leq \dots \leq \nu_N, \quad (1.1)$$

here $\nu_1, \dots, \nu_N$ are the eigenvalues of $\mathcal{V}$, see Sect. 2 for the proof.

It is well known (see [DS] p.1486-1494, [Ge]) that the spectrum $\sigma(\mathcal{K})$ of $\mathcal{K}$ is absolutely continuous and consists of non-degenerated intervals $\sigma_n, n \in \mathbb{Z}$. These intervals are separated by the gaps $g_n = (z_n^-, z_n^+)$ with the length $|g_n| > 0, N_g^- < n < N_g^+$, where $-\infty \leq N_g^- < N_g^+ \leq \infty$ and $N_g = N_g^+ - N_g^- - 1$ is a total number of the gaps.

Introduce the fundamental $2N \times 2N$ -matrix solutions $\psi(t, z)$ of the equation

$$-iJ_1\psi' + V_t\psi = z\psi, \quad z \in \mathbb{C}, \quad \psi(0, z) = I_{2N}, \quad (1.2)$$

and the monodromy $2N \times 2N$ -matrix $\psi(1, z)$. The matrix valued function $\psi(1, \cdot)$ is entire. An eigenvalue of $\psi(1, z)$ is called a *multiplier* of $\mathcal{K}$: to each of them corresponds a solution f of $-iJ_1f' + V_tf = zf$ with $f(t+1) = \tau(z)f(t), t \in [0, 1)$. They are roots of the algebraic equation $D(\tau, z) = 0, \tau, z \in \mathbb{C}$, where $D(\tau, z) = \det(\psi(1, z) - \tau I_N)$, $\tau, z \in \mathbb{C}$. Zeros of the function $\det(\mathcal{M}(z) - I_N)$ (or $\det(\mathcal{M}(z) - I_N)$) are periodic (or anti-periodic) eigenvalues.

There exist many papers about the first order periodic systems $N \geq 2$: Gel'fand and Lidskii [GL], Gohbert and Krein [GK], Krein [Kr], Potapov [Po], [YS] and we mention new papers of Gesztesy and coauthors [CG], [CHGL], [GKM]. The basic results for direct spectral theory for the matrix case were obtained by Lyapunov and Poincaré (see [GL], [Kr], [YS]).

Theorem (Lyapunov, Poincaré). *For each $(V, z) \in \mathcal{H} \times \mathbb{C}$ the matrix-function $\psi(1, z)$ satisfies:*

$$\psi^{-1}(1, \cdot) = -J\psi^\top(1, \cdot)J, \quad (1.3)$$

$$D(\tau, \cdot) = \tau^{2N}D(\tau^{-1}, \cdot), \quad \text{any } \tau \neq 0, \quad \text{where } D(\tau, \cdot) = \det(\psi(1, \cdot) - \tau I_{2N}). \quad (1.4)$$

$$\sigma(\mathcal{K}) = \{z \in \mathbb{C} : |\tau(z)| = 1 \text{ for some multiplier } \tau(z) \text{ of } \psi(1, z)\}. \quad (1.5)$$

If for some $z \in \mathbb{C}$ (or $z \in \mathbb{R}$) $\tau(z)$ is a multiplier of multiplicity $d \geq 1$, then $\tau^{-1}(z)$ (or $\bar{\tau}(z)$) is a multiplier of multiplicity d . Moreover, each $\psi(1, z), z \in \mathbb{C}$, has exactly $2N$ multipliers $\tau_j^{\pm 1}(z), j = 1, \dots, N$. If $\tau(z)$ is a simple multiplier and $|\tau(z)| = 1$, then $\tau'(z) \neq 0$.

The eigenvalues of $\psi(1, z)$ are the zeros of the equation $D(\tau, z) = 0$. This is an algebraic equation in τ of degree $2N$ with coefficients, which are entire in $z \in \mathbb{C}$. It is well known (see e.g. [Fo]) that the zeros $\tau_j(z), j = 1, \dots, 2N$ of $D(\tau, z) = 0$ are (some branches of) analytic functions of z with only algebraic singularities: the zeros $\tau_j(z), j = 1, \dots, 2N$ constitute one or several branches of one or several analytic functions that have only algebraic singularities in $\mathbb{C}$. Thus the number of eigenvalues of $\psi(1, z)$ is a constant N_e with the exception of some special values of z (see below the definition of a resonance). In general, there is a infinite number of such points on the plane. If the functions $\tau_j(z), j = 1, \dots, 2N$ are all distinct, then $N_e = 2N$. If some of them are identical, then we get $N_e < 2N$ and $\psi(1, z)$ is permanently degenerate.

Introduce the matrix-valued function $\mathcal{L}(z) = \frac{1}{2}(\psi(1, z) + \psi^{-1}(1, z)), z \in \mathbb{C}$ and the function $\Phi(z, \nu) = \det(\mathcal{L}(z) - \nu I_{2N}), z, \nu \in \mathbb{C}$. Each zero of $\Phi(\nu, z)$ has multiplicity ≥ 2

and define the Lyapunov function by $\Delta_j(z) = \frac{1}{2}(\tau_j(z) + \tau_j^{-1}(z))$, $j = 1, \dots, N$. The Riemann surface for the multipliers $\tau_j(z)$, $j \in \mathbb{N}_N = \{1, \dots, N\}$ has $2N$ sheets, see (1.4). If $N = 1$, then it has 2 sheets and the Lyapunov function is entire. Similarly, in the case $N \geq 2$ it is more convenient for us to construct the Riemann surface for the Lyapunov function, which has N sheets. We need the following results from [K4].

Theorem 1.1. *Let $V \in \mathcal{H}$. Then there exists an analytic function $\tilde{\Delta}_s$, $s = 1, \dots, N_0 \leq N$ on the N_s -sheeted Riemann surface $\mathcal{R}_s$, $N_s \geq 1$ having the following properties:*

i) There exist disjoint subsets ω_s , $s = 1, \dots, N_0$, $\bigcup \omega_s = \mathbb{N}_N$ such that all branches of $\tilde{\Delta}_s$, $s = 1, 2, \dots, N_0$ has the form $\Delta_j(z) = \frac{1}{2}(\tau_j(z) + \tau_j^{-1}(z))$, $j \in \omega_s$. Moreover, for any $z, \tau \in \mathbb{C}$ the following relations hold true:

$$\det(\mathcal{L}(z) - \nu I_{2N}) = \prod_{s=1}^{N_0} \Phi_s^2(\nu, z), \quad \Phi_s(\nu, z) = \prod_{j \in \omega_s} (\nu - \Delta_j(z)), \quad z, \nu \in \mathbb{C}, \quad (1.6)$$

$$\Delta_j(z) = \cos z + o(e^{|\operatorname{Im} z|}) \quad \text{as} \quad |z| \rightarrow \infty, \quad (1.7)$$

where the functions $\Phi_s(\nu, z)$ are entire with respect to $\nu, z \in \mathbb{C}$. Moreover, if $\Delta_i = \Delta_j$ for some $i \in \omega_k, j \in \omega_s$, then $\Phi_k = \Phi_s$ and $\tilde{\Delta}_k = \tilde{\Delta}_s$.

ii) (The monotonicity property). Let some Δ_j , $j = 1, \dots, N$, be real analytic on some interval $Y = (\alpha, \beta) \subset \mathbb{R}$ and $-1 < \Delta_j(z) < 1$ for any $z \in Y$. Then $\Delta'_j(z) \neq 0$ for each $z \in Y$.

iii) The functions ρ, ρ_s given by (1.8) are entire,

$$\rho = \prod_{s=1}^{N_0} \rho_s, \quad \rho_s(\cdot) = \prod_{i < j, i, j \in \omega_s} (\Delta_i(\cdot) - \Delta_j(\cdot))^2. \quad (1.8)$$

iv) The following identity holds true

$$\sigma(\mathcal{K}) = \mathbb{R} \setminus \bigcup_{j=1}^N g_n, \quad g_n = (z_n^-, z_n^+), \quad N_g^- < n < N_g^+ \quad (1.9)$$

where each gap $g_n = (z_n^-, z_n^+)$ is a bounded interval and $z_n^\pm$ are either periodic (anti-periodic) eigenvalues or real branch points of Δ_j (for some $j = 1, \dots, N$) which are zero of ρ (below we call such point a resonance).

Remark. 1) In the case of 2×2 system the monodromy matrix has exactly 2 eigenvalues τ, τ^{-1} . The Lyapunov function $\frac{1}{2}(\tau + \tau^{-1})$ is an entire function of the spectral parameter. It defines the band-gap structure of the spectrum. By Theorem 1.1, the Lyapunov function for $2N \times 2N$ -matrix operator $\mathcal{K}$ also defines the band-gap structure of the spectrum, but it is the N -sheeted analytic function.

2) We have the following asymptotics (see Sect. 3)

$$\Delta_j(z) = \cos z + \frac{\sin z}{2z} \nu_j + O\left(\frac{e^{|\operatorname{Im} z|}}{z^2}\right), \quad \text{if } V' \in \mathcal{H} \quad (1.10)$$

as $|z| \rightarrow \infty$, $j \in \mathbb{N}_N$. Then firstly, ρ is not a polynomial since ρ is bounded on $\mathbb{R}$. Secondly, if $\nu_{j'} \neq \nu_j$, $j' \neq j$, then (1.10) implies $\Delta_{j'} \neq \Delta_j$.

3) In the case $N = 2$ we determine Δ_1, Δ_2, ρ in terms of the traces of the monodromy matrix. Using (1.4), we have $D(\tau, \cdot) = \tau^4 - T_1\tau^3 + \frac{1}{2}(T_1^2 - T_2)\tau^2 - T_1\tau + 1$, which yields

$$D(\tau, \cdot) = \left(\tau^2 - 2\Delta_1\tau + 1\right)\left(\tau^2 - 2\Delta_2\tau + 1\right), \quad \Delta_1 = \frac{T_1}{2} + \frac{\sqrt{\rho}}{2}, \quad \Delta_2 = \frac{T_1}{2} - \frac{\sqrt{\rho}}{2}, \quad (1.11)$$

see [BBK], where $\rho = \frac{T_2+4}{2} - \frac{T_1^2}{4}$ and $T_m = \text{Tr } \psi(m, z)$, $m = 1, 2$.

Definition. A zero $z_0 \in \mathbb{C}$ of ρ given by (1.8) is a **resonance** of $\mathcal{K}$.

The main goal of this paper is to describe the spectrum of $\mathcal{K}$ and to determine the asymptotics of gaps and resonances, periodic and anti-periodic eigenvalues at high energy. We show that all resonances are real at high energy. Moreover, we prove the existence of complex resonances for some specific periodic potential. We have to underline that in the case of large N the resonances create the gaps in the spectrum of periodic operators, see Theorem 1.2 and remark after Theorem 1.3. If $N = 1$, then 2-periodic eigenvalues create the gaps in the spectrum. In the present paper we use some techniques from [K4] and [BBK], [BK], [CK].

The periodic eigenvalues (n is even) satisfy

$$.. \leq z_N^{-2,+} \leq \underbrace{z_1^{0,+} \leq z_2^{0,+} \leq \dots \leq z_N^{0,+}}_{n=0} \leq \underbrace{z_1^{2,-} \leq z_1^{2,+} \leq \dots \leq z_N^{2,-} \leq z_N^{2,+}}_{n=2} \leq z_1^{4,-} \leq \dots, \quad (1.12)$$

the anti-periodic eigenvalues (n is odd) satisfy

$$.. \leq z_N^{-1,+} \leq \underbrace{z_1^{1,-} \leq z_1^{1,+} \leq \dots \leq z_N^{1,-} \leq z_N^{1,+}}_{n=1} \leq \underbrace{z_1^{3,-} \leq z_1^{3,+} \leq \dots \leq z_N^{3,-} \leq z_N^{3,+}}_{n=3} \leq z_1^{5,-} \leq \dots \quad (1.13)$$

and they have asymptotics

$$z_j^{n,\pm} = \pi n + o(1) \quad \text{as } n \rightarrow \pm\infty, \quad j \in \mathbb{N}_N = \{1, 2, \dots, N\}. \quad (1.14)$$

If $V = 0$, then these eigenvalues have the form $z_j^{n,\pm} = \pi n$, $(n, j) \in \mathbb{Z} \times \mathbb{N}_N$.

Let $\widehat{V}'_n = \int_0^1 V'_t e^{i2\pi n t J_1} dt$, We formulate our first main result

Theorem 1.2. Let $V, V' \in \mathcal{H}$ and let $\zeta_j^{n,\pm}, j \in \mathbb{N}_N$ be eigenvalues of the matrix $\mathcal{V} - iJ_1 \widehat{V}'_n$. Then the periodic and anti-periodic eigenvalues have the following asymptotics:

$$z_j^{n,\pm} = \pi n + \frac{\zeta_j^{n,\pm}}{2\pi n} + O(n^{-2}), \quad j \in \mathbb{N}_N \quad \text{as } n \rightarrow \pm\infty. \quad (1.15)$$

Assume that $\nu_j \neq \nu_{j'}$ for all $j \neq j' \in \omega_s$ for some $s = 1, \dots, N_0$. Then the function ρ_s has the zeros $z_\alpha^{n,\pm}, \alpha = (j, j'), j < j', j, j' \in \omega_s, n \in \mathbb{Z}$, which are real at large $|n|$ and satisfy

$$z_\alpha^{n,\pm} = \pi n + \frac{\nu_j + \nu_{j'}}{4\pi n} + O\left(\frac{|\widehat{V}'_n|}{n} + \frac{1}{n^2}\right), \quad \alpha = (j, j') \quad \text{as } n \rightarrow \pm\infty. \quad (1.16)$$

Let in addition $\nu_1 < \dots < \nu_N$. Then for each $s = 1, \dots, N_0$ and for large $n \rightarrow \pm\infty$ there exists a system of real intervals (gaps) $g_\alpha^n = (z_\alpha^{n-}, z_\alpha^{n+})$ such that

$$z_{j,j'}^{n\pm} = z_{j',j}^{n\pm}, \quad \alpha = (j, j'), \quad j, j' \in \omega_s, \quad z_{j,j_1}^{n-} \leq z_{j,j_1}^{n+} < z_{j,j_2}^{n-} \leq z_{j,j_2}^{n+} < \dots < z_{j,j_{N_s}}^{n-} \leq z_{j,j_{N_s}}^{n+},$$

$$(-1)^n \Delta_j(z) > 1, z \in g_{j,j}^n, \quad \text{and} \quad \overline{\Delta}_j(z) = \Delta_{j'}(z), z \in g_{j,j'}^n, \quad \text{if } j \neq j' \quad (1.17)$$

i) Each branch Δ_j is real and is analytic on the set $(\pi n - \frac{\pi}{2}, \pi n + \frac{\pi}{2}) \setminus \cup_{p \neq j} g_{p,j}^n$ and is not real on $\cup_{p \neq j} g_{p,j}^n$.

ii) If $z_\alpha^{n-} \neq z_\alpha^{n+}$ for some $\alpha = (j, j'), j \neq j'$, then $z_\alpha^{n\pm}$ is the simple branch point (resonance) for the functions $\Delta_j, \Delta_{j'}$. If $z_\alpha^{n-} = z_\alpha^{n+}$, then $\Delta_j, \Delta_{j'}$ are analytic at $z_\alpha^{n\pm}$.

iii) The following asymptotics hold true:

$$z_\alpha^{n\pm} = \pi n + \frac{\nu_j + \nu_{j'} \pm |\widehat{v'_{n,\alpha}}|}{2\pi n} + O\left(\frac{|\widehat{v'_{n,\alpha}}|}{n} + \frac{1}{n}\right), \quad \widehat{v'_{n,\alpha}} = \int_0^1 v'_\alpha(t) e^{-i2\pi n t} dt, \quad \alpha = (j, j'). \quad (1.18)$$

Remark. 1) $N_g = N_g^+ - N_g^- - 1$ is the total number of gaps in the spectrum of $\mathcal{K}$.
2) If $\nu_1 < \dots < \nu_N$, then there exist infinite number of resonances $z_\alpha^{n\pm}, \alpha = (j, j'), j \neq j'$ which form the gaps in the spectrum of $\mathcal{K}$, see below Theorem 1.3. Roughly speaking, resonances "form" the gaps, the number of periodic and anti-periodic eigenvalues is less than the number of resonances. Thus there exists big difference between $N = 1$ and large N . In the first case the endpoints of the gaps are 2-periodic eigenvalues. In the second case, roughly speaking, the endpoints of the gaps are resonances.

In the second main result we describe finite band potentials.

Theorem 1.3. Let $V, V' \in \mathcal{H}$ and let $\nu_1 < \dots < \nu_N$.

(i) If the identity $\nu_1 + \nu_N = \nu_2 + \nu_{N-1} = \dots = \nu_N + \nu_1$ is not fulfilled, then $|N_g^\pm| < \infty$.

(ii) If $\nu_1 + \nu_N = \dots = \nu_N + \nu_1$ holds true and there exists a sequence of indicies $n_k \rightarrow \pm\infty$ such that $|\widehat{v'_{n,\alpha}}|^2 + |n_k|^{-1} = o(|\widehat{v'_{n,\alpha}}|)$ as $k \rightarrow \pm\infty$, for each $\alpha = (j, N+1-j), j \in \mathbb{N}_N$, then $N_g^\pm = \pm\infty$.

Remark. 1) Consider $v = \text{diag}\{v_1, v_2, \dots, v_N\}$, i.e., the case when "variables are separated". The transformation $\mathcal{U}_0 : y = (y_1, \dots, y_{2N})^\top \rightarrow (y_1, y_{N+1}, y_2, y_{N+2}, \dots, y_N, y_{2N})^\top$ gives

$$\mathcal{U}_0 \mathcal{K} \mathcal{U}_0^* = \oplus_1^N \mathcal{K}_j, \quad \mathcal{K}_j = -i\mathbf{j}_1 \frac{d}{dt} + V_j, \quad \mathbf{j}_1 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad V_j = \begin{pmatrix} 0 & v_j \\ v_j^* & 0 \end{pmatrix}.$$

The operator $\mathcal{K}_j$ for the case $N = 1$ is well studied [YS], [K1-3], [Mi]. We have $\nu_j = \int_0^1 |v_j(t)|^2 dt \geq 0$. If $\nu_j < \nu_p$ for some $j < p$, then the number of gaps is $N_g < \infty$.

2) Note that the condition $|\widehat{v'_{n,\alpha}}|^2 + |n|^{-1} = o(|\widehat{v'_{n,\alpha}}|)$, $\alpha = (j, N+1-j), j \in \mathbb{N}_N$ as $n \rightarrow \pm\infty$, holds true for "generic" potentials $V, V' \in \mathcal{H}$. This yields the existence of the real resonance gaps $(z_\alpha^{n-}, z_\alpha^{n+})$ at high energy. The coefficients $\widehat{v'_{n,\alpha}}, \alpha = (j, N+1-j), j \in \mathbb{N}_N$ (the second diagonal of the matrix v) "create" the gaps.

Example of complex resonances. Consider the operator $\mathcal{K}_{\nu,\tau} = -iJ_1 \frac{d}{dt} + V_{\nu,\tau}, \nu = 1, \frac{1}{2}, \frac{1}{3}, \dots, \tau \in \mathbb{R}$ acting in $L^2(\mathbb{R})^4$, where the real periodic potential $V_{\nu,\tau}$ is given by

$$V_{\nu,\tau} = \begin{pmatrix} 0 & v_{\nu,\tau} \\ v_{\nu,\tau} & 0 \end{pmatrix}, \quad v_{\nu,\tau} = - \begin{pmatrix} a & \tau b_\nu \\ \tau b_\nu & 0 \end{pmatrix}, \quad \frac{a}{2\pi} \in \mathbb{R}_+ \setminus \mathbb{N}, \quad b_\nu \in C(\mathbb{T}), \quad (1.19)$$

$$\int_0^1 |b_\nu(t)| dt = 1, \quad \int_0^m b_\nu(t) f(t) dt \rightarrow \int_0^m \delta_{per}(t) f(t) dt, \quad \delta_{per}(t) \equiv \sum_{-\infty}^{\infty} \delta(t - n - \frac{1}{2}), \quad (1.20)$$

as $\nu \rightarrow 0$, for any $f \in C(0, m)$, $m \in \mathbb{N}$.

If $\tau = 0$, then the operator $\mathcal{K}_{\nu,0} = -iJ_1 \frac{d}{dt} + V_{\nu,0}$ has the constant potential $V_{\nu,0}$. In this case there are no gaps in the spectrum of $\mathcal{K}_{\nu,0}$ and all resonances are given by $r_n^0 = \pi n + \frac{a^2}{4\pi n}$, $n \in \mathbb{Z} \setminus \{0\}$ with multiplicity 2. We show that there exist the non-degenerated resonance gaps for small τ, ν . In this example some resonances are real and some are complex.

Proposition 1.4. *Let a potential $V_{\tau,\nu}$ satisfy (1.19), (1.20). Then for each large integer $n_0 \geq 1 + a$ there exist sufficiently small $\nu, \varepsilon > 0$ such that the following statements hold true:*
i) Each function $\rho(z, V_{\tau,\nu})$, $\tau \in (-\varepsilon, \varepsilon)$ in the disk $z \in \mathbb{D}_{n_0} = \{|z| < \pi n_0 + 1\}$ has exactly $4n_0$ simple zeros $r_{n,\nu}^\pm(\tau)$, $1 \leq \pm n \leq n_0$, where $r_{n,\nu}^\pm(\tau)$ is analytic function in the disk $\tau \in \{|\tau| < \varepsilon\}$ and $r_{n,\nu}^\pm(0) = r_n^0$ and the following estimates hold

$$r_{n,\nu}^\pm(\tau) = r_n^0 \pm \tau(\sqrt{R_n} + o(1)) \quad \text{as } \tau \rightarrow 0, \quad \text{where} \quad \begin{cases} R_n < 0 & \text{if } |n| < \frac{a}{2\pi} \\ R_n > 0 & \text{if } |n| > \frac{a}{2\pi} \end{cases}, \quad (1.21)$$

for some constant R_n and if $\tau \in (-\varepsilon, \varepsilon)$, $|n| > \frac{a}{2\pi}$, then $(r_{n,\nu}^-(\tau), r_{n,\nu}^+(\tau)) \subset \mathbb{R}$ is a gap.

ii) Each function $D(1, z, V_{\tau,\nu})D(-1, z, V_{\tau,\nu})$ in the disk $\mathbb{D}_{n_0}$ has exactly $4n_0 + 4$ zeros $z_{n,m}^\pm(\tau, \nu)$, $-n_0 \leq n \leq n_0$, $m = 1, 2$ and the asymptotics $z_{n,m}^\pm(\tau, \nu) = z_{n,m}^{0,\pm} + o(1)$ holds as $\tau \rightarrow 0$.

Remark. 1) If $0 < a < 2\pi$, then $\rho(\cdot, V_{\tau,\nu})$ has only real roots $r_{n,\nu}^\pm(\tau)$ in each large disk $\mathbb{D}_{n_0}$ for sufficiently small τ, ν . If $1 < \frac{a}{2\pi}$, then $\rho(z, V_{\tau,\nu})$ has at least two non-real roots $r_{1,\nu}^\pm(\tau)$ for small τ, ν . 2) We show that operator $\mathcal{K}_{\tau,\nu}$ has new gaps (so-called resonance gaps). The endpoints of the resonance gap are the branch points of the Lyapunov function, and, in general, they are not the periodic (or anti-periodic) eigenvalues. These endpoints are not stable. If they are real (see (1.21)), then we have a gap. If they are complex ($0 < n \leq \frac{a}{2\pi}$), then we have not a gap, we have only the branch points of the Lyapunov function in the complex plane. 3) We have a similar complicated distribution of other resonances, which are poles of S-matrix for scattering for Schrödinger operator with compactly supported potentials on the real line see [K5], [Z].

We consider the conformal mapping associated with the operator $\mathcal{K}$. Introduce the simple conformal mapping $\eta : \mathbb{C} \setminus [-1, 1] \rightarrow \{\zeta \in \mathbb{C} : |\zeta| > 1\}$ by

$$\eta(z) = z + \sqrt{z^2 - 1}, \quad z \in \mathbb{C} \setminus [-1, 1], \quad \text{and} \quad \eta(z) = 2z + o(1) \quad \text{as } |z| \rightarrow \infty. \quad (1.22)$$

Note that $\eta(z) = \bar{\eta}(\bar{z})$, $z \in \mathbb{C} \setminus [-1, 1]$ since $\eta(z) > 1$ for any $z > 1$. Due to the properties of the Lyapunov functions we have $|\eta(\tilde{\Delta}_s(\zeta))| > 1$, $\zeta \in \mathcal{R}_s^+ = \{\zeta \in \mathcal{R}_s : \text{Im } \zeta > 0\}$. Thus we can introduce the quasimomentum k_j , $j = 1, 2, \dots, N$ (we fix some branch of arccos and $\Delta_j(z)$) and the function q_j by

$$k_j(z) = \arccos \Delta_j(z) = i \log \eta(\Delta_j(z)), \quad q_j(z) = \text{Im } k_j(z) = \log |\eta(\Delta_j(z))|, \quad (1.23)$$

$z \in \mathcal{R}_0^+ = \mathbb{C}_+ \setminus \beta_+$, $\beta_+ = \bigcup_{\beta \in \mathcal{B}_\Delta \cap \mathbb{C}_+} [\beta, \beta + i\infty)$ where $\mathcal{B}_\Delta$ is the set of all branch points of the function Δ . The branch points of k_m belong to $\mathcal{B}_\Delta$. Define the **averaged quasimomentum**

k , the **density** p and the **Lyapunov exponent** q by

$$k(z) = p(z) + iq(z) = \frac{1}{N} \sum_1^N k_j(z), \quad q(z) = \text{Im } k(z), \quad z \in \mathcal{R}_0^+. \quad (1.24)$$

For the function $k(z) = p(z) + iq(z)$, $z \in \overline{\mathbb{C}}_+$ we introduce formally integrals

$$Q_n = \frac{1}{\pi} \int_{\mathbb{R}} t^n q(t) dt, \quad I_n^S = \frac{1}{\pi} \int_{\mathbb{R}} t^n q(t) dp(t), \quad I_n^D = \frac{1}{\pi} \iint_{\mathbb{C}_+} |k'_{(n)}(z)|^2 dx dy, \quad (1.25)$$

$n = 0, 1, 2$, here and below $k_{(0)} = k - z$, $k_{(1)} = zk_{(0)}$, $z = x + iy \in \mathbb{C}$. Let C_{us} denote the class of all real upper semi-continuous functions $h : \mathbb{R} \rightarrow \mathbb{R}$. With any $h \in C_{us}$ we associate the "upper" domain $\mathbb{K}(h) = \{k = p + iq \in \mathbb{C} : q > h(p), p \in \mathbb{R}\}$. We formulate our last result.

Theorem 1.5. *i) Let $V \in \mathcal{H}$. Then the averaged quasimomentum $k = \frac{1}{N} \sum_1^N k_j$ is analytic in $\mathbb{C}_+$ and $k : \mathbb{C}_+ \rightarrow \mathbb{K}(\mathbb{C}_+) = \mathbb{K}(h)$ is a conformal mapping for some $h \in C_{us}$. Furthermore, for some branches k_j , $j = 1, \dots, N$ the following asymptotics, identities and estimates hold true:*

$$k(z) - z = -\frac{Q_0 + o(1)}{z} \quad \text{as } |z| \rightarrow \infty, \quad \text{if } y > r|x|, \quad \text{for any } r > 0, \quad (1.26)$$

$$Q_0 = I_0^D + I_0^S = \frac{\|V\|^2}{4N}, \quad (1.27)$$

$$q|_{\sigma_{(N)}} = 0, \quad 0 < q^2|_{\sigma_{(1)} \cup g} \leq 2Q_0, \quad \text{where } \sigma_{(N)} = \{z \in \mathbb{R} : \Delta_1(z), \dots, \Delta_N(z) \in [-1, 1]\},$$

$$\sigma_{(1)} = \{z \in \mathbb{R} : \Delta_m(z) \in [-1, 1], \Delta_p(z) \notin [-1, 1] \text{ some } m, p = 1, \dots, N\},$$

$$\sum_n |g_n|^2 \leq \frac{4\|V\|^2}{N}. \quad (1.28)$$

ii) Let additionally $V' \in \mathcal{H}$. Then the following asymptotics, identities hold true:

$$k(z) - z = -\frac{Q_0}{z} - \frac{Q_1}{z^2} - \frac{Q_2 + o(1)}{z^3} \quad \text{as } |z| \rightarrow \infty, \quad \text{if } y > r|x|, \quad \text{for any } r > 0, \quad (1.29)$$

$$Q_1 = \frac{\text{Tr}}{8N} \int_0^1 -iJ_1 V'_t V_t dt, \quad Q_2 = I_1^D + I_2^S - \frac{Q_0^2}{2} = \frac{\text{Tr}}{16N} \int_0^1 (V_t'^2 + V_t^4) dt. \quad (1.30)$$

A priori estimates for various parameters of the Dirac operator (the norm of a periodic potential, effective masses, gap lengths, height of slits, action variables for NLS and so on) were obtained in [KK1-2], [K1-3], [Mi] only for the case $N = 1$. In order to get the required estimates the authors of [KK1-2], [K1-3], [Mi],... used the global quasi-momentum as the conformal mapping, which was introduced into the spectral theory of the Hill operator by Marchenko-Ostrovski [MO].

The mapping $k : \mathbb{C}_+ \rightarrow \mathbb{K}(h)$ is illustrated in Figure 1 for the case $N = 2$. The integral $I_0^S \geq 0$ is the area between the boundary of $\mathbb{K}(h)$ and the real line. In Figure 1 the upper picture is a domain $\mathbb{K}(h)$ and the points $\tilde{A} = k(A)$, $\tilde{B} = k(B)$,... . The spectral interval

(A, B) (with multiplicity 2) of the z -domain is mapped on the curve $(\tilde{A}, \tilde{B})$ of the k -domain, the interval (a gap) (B, C) of the z -domain is mapped on the vertical slits, which lies on the line $\operatorname{Re} k = 0$. The spectral interval (C, D) (with multiplicity 2) of the z -domain is mapped on the curve $(\tilde{C}, \tilde{D})$ of the k -domain. The spectral interval (D, E) (with multiplicity 4) of the z -domain is mapped on the interval $(\tilde{D}, \tilde{E})$ of the k -domain. The case of the interval (E, J) is similar. The resonance gap (K, L) of the z -domain is mapped on the vertical slits, which lies on the line $\operatorname{Re} k = 2\pi$. In fact we have the graph of the function $h(p), p \in \mathbb{R}$, which coincides with the boundary of $\mathbb{K}(h)$.




Figure 1: The domain $\mathbb{K}(h) = k(\mathbb{C}_+)$ and the graph of the function h

We describe the plan of our paper. In Sect. 2 we obtain the basic properties of fundamental solution $\psi(t, z)$. In Sect. 3 we determine the asymptotics of the fundamental solution $\psi(t, z)$ for the case $V, V' \in \mathcal{H}$. In Sect. 4 we determine the asymptotics of the Lyapunov function and multipliers at high energy and prove Theorem 1.2, 1.5. In Sect. 5 we prove Proposition 1.4. In Sect. 5 we determine the asymptotics of $\det \mathcal{L}(z)$ as $\operatorname{Im} z \rightarrow \infty$.

2 The fundamental solutions

In this section we study ψ . We begin with some notational convention. A vector $h = \{h_n\}_1^N \in \mathbb{C}^N$ has the Euclidean norm $|h|^2 = \sum_1^N |h_n|^2$, while a $N \times N$ matrix A has the operator norm given by $|A| = \sup_{|h|=1} |Ah|$. Note that $|A|^2 \leq \operatorname{Tr} A^* A$.

Recall the identity: if A is $N \times N$ matrix, then

$$\det(A - \nu I_N) = (-1)^N \sum_0^N a_j \nu^{N-j}, \quad a_0 = 1, \quad a_1 = -\mathcal{A}_1, \quad \phi_2 = -\frac{\mathcal{A}_2 + \mathcal{A}_1 \phi_1}{2}, \quad (2.1)$$

and .., $\phi_j = -\frac{1}{j} \sum_1^j \mathcal{A}_k a_{j-k}, \dots, \quad a_N = \det A, \quad \mathcal{A}_m(z) = \operatorname{Tr} A^m.$

Below we need the identity

$$J_1 V = -V J_1, \quad e^{zJ_1} V = V e^{-zJ_1}, \quad (2.2)$$

for any $z \in \mathbb{C}$. The solution of the equation (1.2) satisfies the integral equation

$$\psi(t, z) = \psi_0(t, z) - i \int_0^t e^{izJ_1(t-s)} J_1 V_s \psi(s, z) ds, \quad \psi_0(t, z) = e^{iztJ_1}, \quad t \geq 0, \quad z \in \mathbb{C}. \quad (2.3)$$

It is clear that Eq. (2.3) has a solution as a power series in V given by

$$\psi(t, z) = \sum_{n \geq 0} \psi_n(t, z), \quad \psi_n(t, z) = -i \int_0^t e^{izJ_1(t-s)} J_1 V_s \psi_{n-1}(s, z) ds, \quad n \geq 1. \quad (2.4)$$

Using (2.2), (2.4) we have

$$\psi_1(t, z) = -i \int_0^t e^{izJ_1(t-s)} J_1 V_s e^{izsJ_1} ds = -i \int_0^t e^{izJ_1(t-2s)} J_1 V_s ds, \quad (2.5)$$

$$\psi_2 = -i \int_0^t e^{izJ_1(t-t_1)} J_1 V_{t_1} \psi_1(t_1, z) dt_1 = \int_0^t dt_1 \int_0^{t_1} e^{izJ_1(t-2t_1+2t_2)} V_{t_1} V_{t_2} dt_2. \quad (2.6)$$

Proceeding by induction, we obtain

$$\psi_{2n}(t, z) = \int_0^t dt_1 \dots \int_0^{t_{2n-1}} e^{izJ_1(t-2t_1+2t_2 \dots + 2t_{2n})} V_{t_1} \dots V_{t_{2n}} dt_{2n}, \quad (2.7)$$

$$\psi_{2n+1}(t, z) = -i \int_0^x dt_1 \dots \int_0^{t_{2n}} e^{izJ_1(t-t_1+2t_2 \dots - 2t_{2n+1})} J_1 V_{t_1} \dots V_{t_{2n+1}} dt_{2n+1}. \quad (2.8)$$

We need the following results from [K4].

Lemma 2.1. *Let $V \in \mathcal{H}$. For each $z \in \mathbb{C}$ there exists a unique solution ψ of Eq. (2.3) given by (2.4) and series (2.4) converge uniformly on bounded subsets of $\mathbb{R} \times \mathbb{C} \times \mathcal{H}$. For each $t \geq 0$ the function $\psi(t, z)$ is entire on $\mathbb{C}$. Moreover, for any $n \geq 0$ and $(t, z) \in [0, \infty) \times \mathbb{C}$ the following estimates and asymptotics hold true:*

$$|\psi_n(t, z)| \leq \frac{e^{|\operatorname{Im} z|t}}{n!} \left(\int_0^t |V_s| ds \right)^n, \quad (2.9)$$

$$\left| \psi(t, z) - \sum_{j=0}^{n-1} \psi_j(t, z) \right| \leq \frac{(\sqrt{t} \|V\|)^n}{n!} e^{t|\operatorname{Im} z| + \int_0^t |V_s| ds}, \quad (2.10)$$

$$\psi(t, z) - e^{iztJ_1} = o(e^{t|\operatorname{Im} z|}) \quad \text{as } |z| \rightarrow \infty, \quad (2.11)$$

uniformly on bounded $t \in \mathbb{R}$. If the sequence $V^\nu \rightarrow V$ weakly in $\mathcal{H}$, as $\nu \rightarrow \infty$, then $\psi(t, z, V^\nu) \rightarrow \psi(t, z, V)$ uniformly on bounded subsets of $\mathbb{R} \times \mathbb{C}$.

Below we need the simple properties of matrices $a, b, c \in \mathcal{A} = \left\{ A = \begin{pmatrix} 0 & a_1 \\ a_2 & 0 \end{pmatrix} : a_1, a_2 \text{ is } N \times N \text{ matrix} \right\}$ given by

$$abc, J_1 a, e^{zJ_1} a \in \mathcal{A}, \quad \text{all } z \in \mathbb{C}, \quad (2.12)$$

$$ab = (-a_1 b_1 + a_2 b_2) + (a_1 b_2 - a_2 b_1) J_1, \quad (2.13)$$

$$\text{Tr } a = 0, \quad \text{Tr } J a^n = 0, \quad n \geq 0. \quad (2.14)$$

For any matrices A, B the following identities hold

$$\text{Tr } AB = \text{Tr } BA, \quad \overline{\text{Tr } A} = \text{Tr } A^*. \quad (2.15)$$

Using (2.6)-(2.8) we define the function

$$T_{j,1}(z, V) = \text{Tr} \int_0^j V_{s_1} \int_0^{s_1} V_{s_2} e^{izJ_1(j-2s_1+2s_2)} ds, \quad (2.16)$$

$$T_{j,n}(z, V) = \text{Tr} \psi_{2n}(j, z) = \text{Tr} \int_0^j \dots \int_0^{t_{2n-1}} e^{izJ_1(j-2s_1+2s_2+\dots+2s_{2n})} V_{s_1} \dots V_{s_{2n}} ds, \quad n \geq 2, \quad (2.17)$$

where $s = (s_1, \dots, s_{2n}) \in \mathbb{R}^{2n}$.

Lemma 2.2. *Let $V \in \mathcal{H}$. The functions $T_j(\cdot, V), j = 1, 2, \dots, N$ are entire on $\mathbb{C}$ and $T(z, V) \in \mathbb{R}$ for all $z \in \mathbb{R}$. Moreover, the function $T(z, V)$ satisfies*

$$T_j(z, V) = T_j(z, -V) = 2N \cos jz + \sum_{n \geq 1} T_{j,n}(z, V), \quad (2.18)$$

$$|T_j(z, V)| \leq 2N e^{j(|\text{Im } z| + \|V\|)}, \quad (2.19)$$

$$T_j(z, V) = 2N \cos jz + o(e^{j|\text{Im } z|}) \quad \text{as } |z| \rightarrow \infty, \quad (2.20)$$

and $\text{Tr} \psi_{2n+1}(t, z) = 0$. Series (2.18) converge uniformly on bounded subsets of $\mathbb{C} \times \mathcal{H}$. If a sequence V^ν converges weakly to V in $\mathcal{H}$ as $\nu \rightarrow \infty$, then $T_j(z, V^\nu) \rightarrow T_j(z, V)$ uniformly on bounded subsets of $\mathbb{C}$.

Proof. By Lemma 2.2, series (2.18) converge uniformly and absolutely on bounded subsets of $\mathbb{C} \times \mathcal{H}$. Each term in (2.18) is an entire function of z , then T_j is an entire function of z, V . Moreover, if the sequence V^ν converges weakly to V in $\mathcal{H}$, as $\nu \rightarrow \infty$, then $T_j(z, V^\nu) \rightarrow T_j(z, V)$ uniformly on bounded subsets of $\mathbb{C}$.

We have $T_j = \text{Tr} \psi(j, z) = \text{Tr} \sum_{n \geq 0} \psi_n(j, z)$ and $\text{Tr} \psi_0(j, z) = 2N \cos jz$ for $j \geq 1$. The estimate (2.9) yields (2.19) and (2.11) gives (2.20).

The first relation in (2.12) yields $V_{t_1} \dots V_{t_{2n+1}} \in \mathcal{A}$ for any $t_1, \dots, t_{2n+1} \in \mathbb{R}$. Then relations in (2.12) give $\text{Tr} e^{izJ_1(t-2t_1+2t_2-\dots-2t_{2n+1})} J_1 V_{t_1} \dots V_{t_{2n+1}} = 0$, for any $t, t_1, \dots, t_{2n+1} \in \mathbb{R}$, which together with (2.11) implies $\text{Tr} \psi_{2n+1}(t, z) = 0$.

Below we will show that $T(z, V) \in \mathbb{R}$ for all $z \in \mathbb{R}$ (see (2.28)). ■

We will show (1.1). Consider the self-adjoint operator $Jy' + \Omega(t)y$, $J = \begin{pmatrix} 0 & I_N \\ -I_N & 0 \end{pmatrix}$, acting on the Hilbert space $\oplus_1^{2N} L^2(\mathbb{R})$, where the real matrix Ω is given by

$$\Omega = \begin{pmatrix} \Omega_1 & \Omega_2 \\ \Omega_2 & -\Omega_1 \end{pmatrix} : \quad \Omega_2 = \Omega_2^*, \quad \Omega_1 = \Omega_1^*, \quad \|\Omega\|^2 = \int_0^1 \text{Tr } \Omega^2(t) dt < \infty. \quad (2.21)$$

Let $M(t, z)$, $t \in \mathbb{R}$ be the fundamental solution of the equation $JM' + \Omega M = zM$, $M(0, z) = I_{2N}$. Note that $M(t, z)$ is real for $z \in \mathbb{R}$.

Define the unitary matrix $\mathcal{U} = \frac{1}{\sqrt{2}}(J_1 + iJ) = \mathcal{U}^*$, $\mathcal{U}^2 = I_{2N}$. Using the identities

$$J_2 = \begin{pmatrix} 0 & I_N \\ I_N & 0 \end{pmatrix}, \quad \mathcal{U} J \mathcal{U} = -iJ_1, \quad \mathcal{U} J_1 \mathcal{U} = iJ, \quad \mathcal{U} J_2 \mathcal{U} = -J_2, \quad \Omega = J_1 \Omega_1 + J_2 \Omega_2, \quad (2.22)$$

we deduce that $M_c = \mathcal{U} M \mathcal{U}$ satisfy the equation $-iJ_1 M_c' + \Omega_c M_c = zM_c$, $M_c(0, z) = I_{2N}$, where Ω_c is given by

$$\Omega_c = \Omega_c^* = \mathcal{U} \Omega \mathcal{U} = \begin{pmatrix} 0 & \omega \\ \omega^* & 0 \end{pmatrix} \in \mathcal{H}, \quad \omega = -\Omega_2 + i\Omega_1 = \omega^\top. \quad (2.23)$$

Thus we obtain

$$\int_0^1 \Omega_c^2(t) dt = \mathcal{V}_1 \oplus \mathcal{V}_2, \quad \mathcal{V}_1 = \int_0^1 \omega(t) \omega^*(t) dt, \quad \mathcal{V}_2 = \int_0^1 \omega^*(t) \omega(t) dt, \quad (2.24)$$

and

$$\mathcal{V}_1 = E \mathcal{V}_0 E^* = \overline{\mathcal{V}_2} \geq 0, \quad \mathcal{V}_0 = \text{diag}\{\nu_1, \dots, \nu_N\}, \quad (2.25)$$

for some unitary matrix E and the diagonal matrix $\mathcal{V}_0$. Define the unitary matrix $\mathcal{E} = E \oplus \overline{E}$. The function $\psi(t, z) = \mathcal{E}^* M_c(t, z) \mathcal{E}$ satisfies $-iJ_1 \psi' + V \psi = z\psi$, $\psi(0, z) = I_{2N}$, where

$$\mathcal{E} = E \oplus \overline{E}, \quad V = \mathcal{E}^* \Omega_c \mathcal{E} = \begin{pmatrix} 0 & v \\ v^* & 0 \end{pmatrix}, \quad v = E^* \omega \overline{E}, \quad (2.26)$$

$$\int_0^1 V_t^2 dt = \mathcal{V}_0 \oplus \mathcal{V}_0, \quad (2.27)$$

$$\text{Tr } \psi(t, z) = \text{Tr } M(t, z), \quad (2.28)$$

which gives $V \in \mathcal{H}$ and $\text{Tr } \psi(1, z) \in \mathbb{R}$ for all $z \in \mathbb{R}$.

It is well known that for real Ω we have $M(t, z) J M(t, z)^\top = J$ (see [GL], [YS]). Then $M_c = \mathcal{U} M \mathcal{U}$ and (2.22) give

$$-iJ_1 = M_c(1, z) (-iJ_1) \mathcal{U} \mathcal{U}^\top (M_c(1, z)^\top) \mathcal{U}^\top \mathcal{U}, \quad (-iJ_1) \mathcal{U} \mathcal{U}^\top = -J,$$

which yields

$$M_c(1, z) J M_c(1, z)^\top = J. \quad (2.29)$$

The similar arguments and $\psi = \mathcal{E}^* M_c \mathcal{E}$ imply

$$\psi \mathcal{E}^* J \mathcal{E}^{*\top} \psi^\top \mathcal{E}^\top = \mathcal{E}^* J, \quad \mathcal{E}^* J \mathcal{E}^{*\top} = J,$$

which yields $\psi J \psi^\top = J$ and (1.3) is proved for $V \in \mathcal{H}$.

3 Estimates of ψ for the case $V' \in \mathcal{H}$

Sect. 2 does not give the needed estimates of the fundamental solution ψ at high energy. In order to determine the asymptotics of ψ we will do some modification. Define the integral operator K and the matrix-valued function $a_t(z)$ by

$$(Kf)(t) = \int_0^t e_{t-s} W_s f(s) ds, \quad W = -iJ_1 V^2 - V', \quad e_t = e_t(z) = e^{itzJ_1}, \quad a_t(z) = I - \frac{V_t}{2z}. \quad (3.1)$$

where $e_t = e_t(z) = e^{itzJ_1}$. Introduce

$$\widehat{V}'(z) = \int_0^1 V'_t e^{i2tzJ_1} dt, \quad \widehat{V}^3(z) = \int_0^1 V_t^3 e^{i2tzJ_1} dt, \quad \mathcal{V} = \int_0^1 V_t^2 dt. \quad (3.2)$$

Lemma 3.1. *For each $(z, V') \in (\mathbb{C} \setminus \{0\}) \times \mathcal{H}$ the solution $\psi = a^{-1}\Psi a_0$, where Ψ satisfies*

$$\Psi = \psi_0 + \varepsilon K a^{-1} \Psi, \quad \Psi = a \psi a_0^{-1}, \quad a_0 = I - \varepsilon V_0, \quad \varepsilon = \frac{1}{2z}, \quad (3.3)$$

$$\Psi = \psi_0 + \sum_{n \geq 1} \Psi_n, \quad \Psi_n = \varepsilon^n (K a^{-1})^n \psi_0, \quad (3.4)$$

where series (3.4) converge uniformly on bounded subsets of $\mathbb{R} \times (\mathbb{C} \setminus \{0\}) \times \mathcal{H}$. Moreover, if $\sup_{t \in \mathbb{R}} |V_t| \leq |z|$, then for any $j-1, m \in \mathbb{N}$ the following estimates are fulfilled:

$$|\Psi_n(t, z)| \leq \frac{e^{|\operatorname{Im} z|t}}{n! |z|^n} \left(\int_0^t |W_s| ds \right)^n, \quad (3.5)$$

$$|\Psi(m, z) - \psi_0(m, z) - \sum_1^{j-1} \Psi_n(m, z)| \leq \frac{\varkappa^j}{j!} e^{m(|\operatorname{Im} z| + \varkappa)}, \quad \varkappa \equiv \frac{\|V\|^2 + \|V'\|}{|z|}, \quad (3.6)$$

$$\Psi_1(1, z) = -\frac{e^{izJ_1}}{2z} \left(iJ_1 \mathcal{V} + \widehat{V}'(z) + \frac{1}{2z} \left(iJ_1 \widehat{V}^3(z) + \int_0^1 V'_t V_t dt \right) \right) + O\left(\frac{e^{|\operatorname{Im} z|}}{z^3}\right), \quad (3.7)$$

$$\Psi_2(1, z) = -\frac{e^{izJ_1}}{4z^2} \int_0^1 dt \int_0^t \left(V_t^2 V_s^2 - iJ_1 V_t^2 V'_s e_{2s} - iV'_t e_{2t} J_1 V_s^2 - V'_t e_{2t} V'_s e_{2s} \right) ds + O\left(\frac{e^{|\operatorname{Im} z|}}{z^3}\right) \quad (3.8)$$

as $|z| \rightarrow \infty$ and where $e_t = e^{itzJ_1}$.

Proof. Using (2.2), (2.3), $\varepsilon = \frac{1}{2z}$ and integrating by parts we get

$$\begin{aligned} \psi(t, z) - e^{iztJ_1} &= -i \int_0^t J_1 e^{izJ_1(t-s)} V_s \psi(s, z) ds = -i \int_0^t e^{izJ_1(t-2s)} J_1 \left(V_s e^{-izsJ_1} \psi(s, z) \right) ds \\ &= \varepsilon e^{izJ_1(t-2s)} \left(V_s e^{-izsJ_1} \psi(s, z) \right) \Big|_0^t - \varepsilon \int_0^t e^{izJ_1(t-2s)} \left(V_s e^{-izsJ_1} \psi(s, z) \right)' ds \\ &= \varepsilon \left(V_t \psi(t, z) - e^{iztJ_1} V_0 \right) - \varepsilon \int_0^t e^{izJ_1(t-2s)} \left(V'_s e^{-izsJ_1} + i e^{izsJ_1} J_1 V_s^2 \right) \psi(s, z) ds. \end{aligned}$$

Thus we obtain

$$a(t, z)\psi(t, z) = e^{iztJ}a(0, z) + \varepsilon \int_0^t e^{izJ(t-s)}W_s\psi(s, z)ds$$

which yields (3.3). We will show (3.4)-(3.6). Using $|1/a(t, z)| \leq 2$ for $\sup_{t \in \mathbb{R}} |V_t| \leq |z|$ and $|\psi_0(t, z)| \leq e^{|\operatorname{Im} z|t}$, we have

$$\begin{aligned} |\Psi_n(t, z)| &\leq 2|\varepsilon| \int_0^t e^{|\operatorname{Im} z|(t-t_1)} |W_{t_1}| |\Psi_{n-1}(t_1, z)| dt_1 \\ &\leq (2|\varepsilon|)^n \int_0^t dt_1 \int_0^{t_1} dt_2 \dots \int_0^{t_{n-1}} e^{|\operatorname{Im} z|t} |W_{t_1}| \dots |W_{t_n}| dt_1 \leq \frac{(2|\varepsilon|)^n}{n!} e^{|\operatorname{Im} z|t} \left(\int_0^t |W_{t_1}| dt_1 \right)^n, \end{aligned}$$

which gives (3.5). Estimates (3.5) and $\int_0^m |W(t)| dt \leq m(\|V'\| + \|V\|^2)$ imply (3.4) and (3.6). We get

$$\Psi_1(1, z) = \varepsilon K a^{-1} \psi_0 = \varepsilon K \psi_0 + \varepsilon^2 K V \psi_0 + O(\varepsilon^3 e^{|\operatorname{Im} z|}). \quad (3.9)$$

Recall $e_t = e^{iztJ_1}$. (2.10) implies

$$K\psi_0 = - \int_0^1 e_{1-t}(iJ_1 V_t^2 + V_t') e_t dt = -ie_1 J_1 \mathcal{V} - e_1 \widehat{V'}(z),$$

$$KV\psi_0 = - \int_0^1 e_{1-t}(iJ_1 V_t^2 + V_t') V_t e_t dt = -ie_1 J_1 \widehat{V^3}(z) - e_1 \int_0^1 V_t' V_t dt,$$

which yields (3.7). Consider the second term $\Psi_2 = \varepsilon^2 (K a^{-1})^2 \psi_0 = \varepsilon^2 \Psi_{20} + O(\varepsilon^3 e^{|\operatorname{Im} z|})$, where

$$\begin{aligned} \Psi_{20} &= \int_0^1 dt \int_0^t e_{1-t}(iJ_1 V_t^2 + V_t') e_{t-s}(iJ_1 V_s^2 + V_s') e_s ds \\ &= \int_0^1 dt \int_0^t e_{1-t} \left[-J_1 V_t^2 e_{t-s} J_1 V_s^2 + iJ_1 V_t^2 e_{t-s} V_s' - iV_t' e_{t-s} J_1 V_s^2 + V_t' e_{t-s} V_s' \right] e_s \\ &= -e_1 \int_0^1 dt \int_0^t \left[V_t^2 V_s^2 - ie_{-2s} J_1 V_t^2 V_s' - ie_{-2t} V_t' J_1 V_s^2 - e_{-2t+2s} V_t' V_s' \right] ds \end{aligned}$$

which gives (3.8). ■

In order to determine the asymptotics of the Lyapunov function we need the following modification. Substituting $\psi^{-1} = -J\psi^\top J$ into $\Psi(1, z)^{-1} = a_0 \psi(1, z)^{-1} a_0^{-1}$ we get

$$\Psi(1, z)^{-1} = -a_0 J a_0 \Psi(1, z)^\top a_0^{-1} J a_0^{-1} = -c J \Psi(1, z)^\top J c^{-1}, \quad c = I - \varepsilon^2 V_0^2, \varepsilon = \frac{1}{2z},$$

which yields

$$L = \frac{1}{2} \left(\Psi(1, z) + \Psi(1, z)^{-1} \right) = \frac{1}{2} \left(\Psi(1, z) - c J \Psi^\top(1, z) J c^{-1} \right). \quad (3.10)$$

We determine the asymptotics of $\Psi(1, z)$. Using (3.6), we have

$$\begin{aligned}\Psi &= \psi_0 + \varepsilon K a^{-1} \psi_0 + \varepsilon^2 K a^{-1} K a^{-1} \psi_0 + \varepsilon^2 K a^{-1} K a^{-1} K a^{-1} \psi_0 + O(\varepsilon^4 e^{|\operatorname{Im} z|}) \\ &= \psi_0 + \varepsilon K(I + \varepsilon V + \varepsilon^2 V^2) \psi_0 + \varepsilon^2 K(I + \varepsilon V) K(I + \varepsilon V) \psi_0 + \varepsilon^2 K^3 \psi_0 + O(\varepsilon^4 e^{|\operatorname{Im} z|}) \\ &= \psi_0 + \varepsilon \varkappa_1 + \varepsilon^2 \varkappa_2 + \varepsilon^3 \varkappa_3 + O(\varepsilon^4 e^{|\operatorname{Im} z|}),\end{aligned}$$

where

$$\varkappa_1 = K \psi_0, \quad \varkappa_2 = (KV + K^2) \psi_0, \quad \varkappa_3 = (KV^2 + K^2 V + KVK + K^3) \psi_0. \quad (3.11)$$

We get

$$\begin{aligned}2L &= \psi_0 + \varepsilon \varkappa_1 + \varepsilon^2 \varkappa_2 + \varepsilon^3 \varkappa_3 - (I - \varepsilon^2 V_0^2) J(\psi_0 + \varepsilon \varkappa_1 + \varepsilon^2 \varkappa_2 + \varepsilon^3 \varkappa_3)^\top J(I + \varepsilon^2 V_0^2) + O(\varepsilon^4 e^{|\operatorname{Im} z|}) \\ &= (\psi_0 - J\psi_0^\top J) + \varepsilon(\varkappa_1 - J\varkappa_1^\top J) + \varepsilon^2(\varkappa_2 - J\varkappa_2^\top J + V^2 J\psi_0^\top J - J\psi_0^\top J V_0^2) \\ &\quad + \varepsilon^3(\varkappa_3 - J\varkappa_3^\top J + V_0^2 J\varkappa_1^\top J - J\varkappa_1^\top J V_0^2) + O(\varepsilon^4 e^{|\operatorname{Im} z|}).\end{aligned}$$

Thus we have

$$L = \cos z + \varepsilon L_1 + \varepsilon^2 L_2 + \varepsilon^3 L_3 + O(\varepsilon^4 e^{|\operatorname{Im} z|}), \quad (3.12)$$

where

$$L_1 = \frac{\varkappa_1 - J\varkappa_1^\top J}{2}, \quad L_2 = \frac{\varkappa_2 - J\varkappa_2^\top J}{2}, \quad L_3 = \frac{1}{2}(\varkappa_3 - J\varkappa_3^\top J + V_0^2 J\varkappa_1^\top J - J\varkappa_1^\top J V_0^2). \quad (3.13)$$

Below we need the identities

$$J^* V^\top J = -V, \quad J^* W^\top J = -W, \quad J^* J_1 J = -J_1. \quad (3.14)$$

We determine the asymptotics of L . Let $u_t = \int_0^t V_s^2 ds$, $f_t = \int_0^t V_s' e_{2s} ds$.

Lemma 3.2. *If $V, V' \in \mathcal{H}$, then asymptotics (1.10), the following identities and asymptotics are fulfilled:*

$$L_1 = \mathcal{V} \sin z, \quad z \in \mathbb{C}, \quad \text{where } \mathcal{V} = \int_0^1 V_t^2 dt, \quad (3.15)$$

$$L_2 = i \frac{e_1 J_1}{2} (\mathcal{V} \widehat{V'} + \widehat{V'} \mathcal{V}) - i \sin z \int_0^1 J_1 V_t' V_t dt + L_{21} + L_{22}, \quad (3.16)$$

$$\begin{aligned}L_{21} &= -\frac{1}{2} \int_0^1 (e_1 u' u + e_{-1} u u') dt, \quad L_{22} = \frac{1}{2} \int_0^1 (e_1 f' f + e_{-1} f f') dt, \\ L_{22} &= o(e^{|\operatorname{Im} z|}),\end{aligned} \quad (3.17)$$

$$L_2(\pi n) = \frac{(-1)^n}{2} \left(-\mathcal{V}^2 + i J_1 (\mathcal{V} \widehat{V_n'} + \widehat{V_n'} \mathcal{V}) + (\widehat{V_n'})^2 \right), \quad \widehat{V_n'} = \widehat{V'}(\pi n), \quad (3.18)$$

$$\Delta_j(z) = \cos\left(z - \frac{\nu_j}{2z}\right) + \frac{O(|\widehat{V_n'}| + |n|^{-1})}{n^2}, \quad \text{as } z = \pi n + O\left(\frac{1}{n}\right), \quad j = 1, \dots, N. \quad (3.19)$$

Proof. Recall $e_t = e^{iztJ_1}$. (3.11),(4.1) give

$$J^* \varkappa_1^\top J = J^*(K\psi_0)^\top J = J^* \int_0^1 (e_{1-t} W_t e_t)^\top J dt = \int_0^1 e_{-t} J^* W_t^\top J e_{t-1} dt = - \int_0^1 e_{-t} W_t e_{t-1} dt.$$

Then

$$2L_1 = \int_0^1 (e_{1-t} W_t e_t - e_{-t} W_t e_{t-1}) dt = -iJ_1 \int_0^1 (e_{1-t} V_t^2 e_t - e_{-t} V_t^2 e_{t-1}) dt = -iJ_1 \int_0^1 (e_1 - e_{-1}) V_t^2 dt$$

which yields (3.15). We determine $L_2 = \frac{1}{2}(\varkappa_2 - J\varkappa_2^\top J)$. Using (3.11),(4.1) we get

$$\varkappa_2 = (KV + K^2)\psi_0 = \int_0^1 e_{1-t} W_t \left(V_t e_t + \int_0^t (e_{t-s} W_s e_s ds) \right) dt$$

$$J^* \varkappa_2^\top J = J^* \int_0^1 \left(e_t V_t^\top + \int_0^t (e_s W_s^\top e_{s-t} ds) \right) W_t^\top e_{1-t} dt J = \int_0^1 \left(e_{-t} V_t + \int_0^t (e_{-s} W_s e_{s-t} ds) \right) W_t e_{t-1} dt.$$

This yields

$$L_2 = F + S, \quad S = \frac{1}{2} \int_0^1 dt \int_0^t (e_{1-t} W_t e_{t-s} W_s e_s + e_{-s} W_s e_{s-t} W_t e_{t-1}) ds$$

and using $W = -iJ_1 V^2 - V'$ we obtain

$$\begin{aligned} F &= \int_0^1 (e_{1-t} W_t V_t e_t + e_{-t} V_t W_t e_{t-1}) \frac{dt}{2} = - \int_0^1 (e_{1-t} (iJ_1 V_t^2 + V_t') V_t e_t + e_{-t} V_t (iJ_1 V_t^2 + V_t') e_{t-1}) \frac{dt}{2} \\ &= -\frac{1}{2} \int_0^1 (e_1 V_t' V_t + e_{-1} V_t V_t') dt = \frac{e_{-1} - e_1}{2} \int_0^1 V_t' V_t dt = -iJ_1 \sin z \int_0^1 V_t' V_t dt \end{aligned}$$

since $\int_0^1 V_t V_t' dt = - \int_0^1 V_t' V_t dt$. Consider the second term,

$$\begin{aligned} 2S &= \int_0^1 dt \int_0^t (e_{1-t} (iJ_1 V_t^2 + V_t') e_{t-s} (iJ_1 V_s^2 + V_s') e_s + e_{-s} (iJ_1 V_s^2 + V_s') e_{s-t} (iJ_1 V_t^2 + V_t') e_{t-1}) ds \\ &= \int_0^1 dt \int_0^t (e_{1-t} \left[-J_1 V_t^2 e_{t-s} J_1 V_s^2 + iJ_1 V_t^2 e_{t-s} V_s' + iV_t' e_{t-s} J_1 V_s^2 + V_t' e_{t-s} V_s' \right] e_s + \\ &\quad + e_{-s} \left[-J_1 V_s^2 e_{s-t} J_1 V_t^2 + iJ_1 V_s^2 e_{s-t} V_t' + iV_s' e_{s-t} J_1 V_t^2 + V_s' e_{s-t} V_t' \right] e_{t-1} \\ &= \int_0^1 dt \int_0^t \left(\left[-e_1 V_t^2 V_s^2 + ie_{1-2s} J_1 V_t^2 V_s' + ie_{1-2t} V_t' J_1 V_s^2 + e_{1-2t+2s} V_t' V_s' \right] \right. \\ &\quad \left. + \left[-e_{-1} V_s^2 V_t^2 + ie_{1-2t} J_1 V_s^2 V_t' + ie_{1-2s} V_s' J_1 V_t^2 + e_{-1+2t-2s} V_s' V_t' \right] \right) ds \\ &= - \int_0^1 dt (e_1 u' u + e_{-1} u u') dt + \frac{ie_1 J_1}{2} (\mathcal{V} \widehat{V'} + \widehat{V'} \mathcal{V}) + \int_0^1 dt (e_1 f' f + e_{-1} f f') dt \end{aligned}$$

where $\widehat{V}'(z) = \int_0^1 V'(t)e_{2t}dt$ and $u = \int_0^t V_s^2 ds$, $f = \int_0^t V'_s e_{2s} ds$, and here we used

$$\begin{aligned} \int_0^1 dt \int_0^t (e_1 V_t^2 V_s^2 + e_{-1} V_s^2 V_t^2) dt ds &= \int_0^1 dt (e_1 u' u + e_{-1} u u') dt, \\ \int_0^1 (u' f + f' u + u f' + f u') dt &= u_1 f_1 + f_1 u_1 = \mathcal{V} \widehat{V}' + \widehat{V}' \mathcal{V}, \\ \int_0^1 dt \int_0^t (e_{1-2t+2s} V'_t V'_s + e_{2t-2s-1} V'_s V'_t) ds &= \int_0^1 dt (e_1 f' f + e_{-1} f f') dt. \end{aligned}$$

We have $e_{\pm 1} = (-1)^n I_{2N}$ at $z = \pi n$ and then

$$L_{21}(\pi n) = -\frac{(-1)^n}{2} \int_0^1 (u' u + u u') dt = -(-1)^n \frac{\mathcal{V}^2}{2}, \quad (3.20)$$

$$L_{22}(\pi n) = \frac{(-1)^n}{2} \int_0^1 (f' f + f f') dt = \frac{(-1)^n}{2} f^2(\pi n) = \frac{(-1)^n}{2} (\widehat{V}')^2(\pi n), \quad (3.21)$$

which yields (3.18). Using (3.12), (3.15) (3.18) we obtain

$$L(z) = \cos(z - \varepsilon \mathcal{V}) + \varepsilon^2 \left(L_2(\pi n) + (-1)^n \frac{\mathcal{V}^2}{2} \right) + O(\varepsilon^3) \quad \text{as } z = \pi n + O(1/n). \quad (3.22)$$

Recall the simple fact: Let A, B be matrices and $\sigma(B)$ be spectra of B . If A be normal, then $\text{dist}\{\sigma(A), \sigma(A+B)\} \leq |B|$ (see [Ka, p.291]).

The normal operator $\cos(z - \varepsilon \mathcal{V})$ has the eigenvalues $\cos(z - \varepsilon \nu_j)$, $j = 1, \dots, N$ with the multiplicity 2. Using the result from [Ka] and asymptotics (3.12) and identity (3.18) we deduce that the eigenvalues $\Delta_j(z)$ of matrix $L(z)$ satisfy the asymptotics (3.19). The proof of (1.10) is similar. ■

4 Proof of the main theorems

We need the following results from [K4].

Lemma 4.1. *Let $V, V' \in \mathcal{H}$. Then the following asymptotics hold true:*

$$\Phi(z, \nu) = (\cos z - \nu)^{2N} + o(e^{N|\text{Im } z|}) \quad \text{as } |z| \rightarrow \infty, \quad (4.1)$$

where $|\nu| \leq A_0$ for some constant $A_0 > 0$. Moreover, there exists an integer n_0 such that:

i) the function $\Phi(z, 1)$ has exactly $N(2n_0 + 1)$ roots, counted with multiplicity, in the disc $\{|z| < \pi(2n_0 + 1)\}$ and for each $|n| > n_0$, exactly $2N$ roots, counted with multiplicity, in the domain $\{|z - 2\pi n| < \frac{\pi}{2}\}$. There are no other roots.

ii) the function $\Phi(z, -1)$ has exactly $2Nn_0$ roots, counted with multiplicity, in the disc $\{|z| < 2\pi n_0\}$ and for each $|n| > n_0$, exactly $2N$ roots, counted with multiplicity, in the domain $\{|z - \pi(2n + 1)| < \frac{\pi}{2}\}$. There are no other roots.

iii) Let in addition $\nu_i \neq \nu_j$ for all $i \neq j \in \omega_s$ for some $s = 1, \dots, N_0$. Then the function ρ_s has exactly $2N_s(N_s - 1)n_0$ roots, counted with multiplicity, in the disc $\{|z| < \pi(n_0 + \frac{1}{2})\}$ and for each $|n| > n_0$, exactly $N_s(N_s - 1)$ roots, counted with multiplicity, in the domain $\{|z - \pi n| < \frac{\pi}{2}\}$. There are no other roots. Moreover,

$$\rho_s(z) = c_s \left(\frac{\sin z + o(e^{|\operatorname{Im} z|})}{2z} \right)^{N_s(N_s-1)}, \quad c_s = \prod_{j,k \in \omega_s} (\nu_j - \nu_k)^2, \quad |z| \rightarrow \infty. \quad (4.2)$$

ii) All zeros of ρ_s are given by $z_\alpha^{a,\pm}$, $\alpha = (j, k)$, $j < k$, $j, k \in \omega_s$ and $n \in \mathbb{Z} \setminus \{0\}$. Furthermore, they satisfy

$$z_\alpha^{n\pm} = \pi n + \frac{\nu_j + \nu_k + o(1)}{2\pi n}, \quad \alpha = (j, k), \quad n \rightarrow \pm\infty. \quad (4.3)$$

Proof of Theorem 1.2. i) We determine asymptotics (1.15) for $z_j^{n\pm}$ as $n \rightarrow \pm\infty$, $j = 1, \dots, N$. Lemma 3.2 yields $|z_j^{n\pm} - \pi n| < \frac{\pi}{2}$, as $n \rightarrow \infty$, $j = 1, 2, \dots, 2N$. Lemma 3.1 gives $\Delta_m(z) = \cos(z - \frac{\nu_m}{2z}) + O(1/z^2)$, $m = 1, \dots, N$ as $z = \pi n + O(1)$. For each $m = 1, \dots, N$ there exists j such that $\Delta_j(z_m^{n\pm}) = (-1)^n$. Thus we have $z_m^{n\pm} = \pi n + O(1/n)$. Define the lokal parameter μ by $z = \pi n + \varepsilon\mu$, $\varepsilon = \frac{1}{2\pi n}$. In order to improve these asymptotics of $z_m^{n\pm}$ we need asymptotics of the $\Psi(1, z)$ as $z = \pi n + O(1/n)$, $n \rightarrow \pm\infty$ given by (3.7)

$$\Psi(1, z) = e^{izJ_1} \left(1 - i\varepsilon J_1 \Gamma_n + O(\varepsilon^2) \right), \quad \Gamma_n = \mathcal{V} - iJ_1 \widehat{V}'_n, \quad \widehat{V}'_n = \int_0^1 V'(s) e^{i2\pi n J_1 s} ds$$

where $\mathcal{V} = \int_0^1 V^2(t) dt = \mathcal{V}_0 \oplus \mathcal{V}_0$, $\mathcal{V}_0 = \operatorname{diag}\{\nu_1, \dots, \nu_N\}$, $0 \leq \nu_1 \leq \nu_2 \leq \dots \leq \nu_N$. Thus we get

$$A = \frac{(-1)^n \Psi(1, z) - I}{-i\varepsilon} = \frac{e^{i\varepsilon\mu J_1} (I - i\varepsilon J_1 \Gamma_n + O(\varepsilon^2)) - I}{-i\varepsilon} = J_1 (\Gamma_n - \mu + O(\varepsilon)). \quad (4.4)$$

Hence we study the zeros of the equation

$$\det(\mathcal{V} - iJ_1 \widehat{V}'_n + O(\varepsilon) - \mu) = 0, \quad \widehat{V}'_n = \begin{pmatrix} 0 & \widehat{v}'_n \\ \widehat{v}'_n^* & 0 \end{pmatrix}, \quad \widehat{v}'_n = \int_0^1 v'(t) e^{-i2\pi n t} dt, \quad (4.5)$$

where $\mu \in \mathbb{C}$. We will use the standard arguments from the perturbation theory (see [Ka,p.291]). Let A, B be bounded operators, A be a normal operator and $\sigma(A), \sigma(B)$ be spectra of A, B . Then $\operatorname{dist}\{\sigma(A), \sigma(B)\} \leq \|A - B\|$.

Let $\zeta_m^{n\pm}$, $(m, n) \in \{1, 2, \dots, N\} \times \mathbb{Z}$ be the eigenvalues of the self-adjoint operator $\mathcal{V} - iJ_1 \widehat{V}'_n$. Using the arguments from the perturbation theory (see [Ka,p.291]), we obtain that Eq. (4.5) has zeros $\omega_m^{n\pm}$, $(m, n) \in \{1, 2, \dots, N\} \times \mathbb{Z}$ such that $\omega_m^{n\pm} = \zeta_m^{n\pm} + O(n^{-1})$ as $n \rightarrow \pm\infty$, which yields (1.15).

Consider the case $\nu_1 < \dots < \nu_N$. We shall determine asymptotics (1.18) for the case $\alpha = (m, m)$, $m = 1, \dots, N$. Let $z_\alpha^{n\pm} = \pi n + \varepsilon\mu$ and $\mu - \nu_m = \xi \rightarrow 0$. Using the simple transformation (unitary), i.e., changing the lines and columns, we obtain

$$\det(\mathcal{V} - iJ_1 \widehat{V}'_n + O(\varepsilon) - \mu) = \det \begin{pmatrix} A_1 & A_2 \\ A_3 & A_4 \end{pmatrix} = \det A_4 \det K,$$

$$A_1 = \begin{pmatrix} -\xi & b \\ \bar{b} & -\xi \end{pmatrix} + O(\varepsilon), \quad b = -i\widehat{v}_{n',\alpha}, \alpha = (m, m), \quad A_2, A_3 = O(\delta_n), \quad \delta_n = |\widehat{V}'_n| + |\varepsilon|,$$

$$A_4 = \mathcal{V}_{0m} \oplus \mathcal{V}_{0m} + O(\delta_n), \quad \mathcal{V}_{0m} = \text{diag}\{\nu_j - \xi, j \neq m\},$$

$$K = A_1 - A_2 A_4^{-1} A_3 + O(\varepsilon) = A_1 + O(\phi), \quad \phi = |\varepsilon| + |\widehat{V}'_n|^2,$$

which yields

$$0 = \det K = \xi^2 - |b|^2 - \xi b_1 + ab_2 + bb_3 + O(\phi^2), \quad b_1, b_2, b_3 = O(\phi),$$

where b_1, b_2, b_3 are analytic functions of ξ . Rewriting the last equation in the form $(\xi + \alpha)^2 = (c + \beta)^2 + O(\phi^2)$, $\alpha, \beta = O(\phi)$ and using the estimate $\sqrt{x^2 + y^2} - x \leq y$ for $x, y \geq 0$ we get $\xi = \pm c + O(\phi)$, which yields (1.18) for $i = j$.

Consider the resonances. We shall determine asymptotics (1.18) for the case $\nu_j \neq \nu_{j'}$ for all $j \neq j' \in \omega_s$ for some $s = 1, \dots, N_0$. By Lemma 3.2, the zeros of ρ_s have the form $z_\alpha^{n\pm}, \alpha = (j, j'), j, j' \in \omega_s, j < j', n \in \mathbb{Z}$ and satisfy $|z_\alpha^{n\pm} - \pi n| < \pi/2$.

Asymptotics (1.10) yields $\Delta_j(z) - \Delta_{j'}(z) = (\nu_j - \nu_{j'}) \frac{\sin z}{2z} + O(z^{-2} e^{|\text{Im} z|})$, $|z| \rightarrow \infty$. Then $|z_\alpha^{n\pm} - \pi n| < \pi/2$ yields $z_\alpha^{n\pm} = \pi n + O(1/n)$ as $n \rightarrow \infty$.

We have the identity $\Delta_j(z) - \Delta_{j'}(z) = 0$ at $z = z_\alpha^{n\pm}$. Then using (3.19) we have

$$\cos\left(z_\alpha^{n\pm} - \frac{\nu_j}{2\pi n}\right) - \cos\left(z_\alpha^{n\pm} - \frac{\nu_{j'}}{2\pi n}\right) = 2(-1)^n \sin \frac{\nu_{j'} - \nu_j}{4\pi n} \sin\left(z_\alpha^{n\pm} - \pi n - \frac{\nu_j + \nu_{j'}}{4\pi n}\right) = \frac{O(\delta_n)}{n^2}$$

which yields (1.16), i.e.,

$$z_\alpha^{n\pm} = \pi n + \varepsilon(a_+ + \widetilde{z}_\alpha^{n\pm}), \quad a_\pm = \frac{\nu_j \pm \nu_{j'}}{2}, \quad \varepsilon = \frac{1}{2\pi n}, \quad \widetilde{z}_\alpha^{n\pm} = O(\delta_n). \quad (4.6)$$

We shall show that for large n in the neighborhood of each $\pi n + \varepsilon a_+$ the function $(\Delta_j(z) - \Delta_{j'}(z))^2$ has two real zeros resonances (counted with multiplicity). Introduce the functions

$$f_m(\mu) = 2(2\pi n)^2(1 - (-1)^n \Delta_m(\pi n + \varepsilon \mu)) = (\mu - \nu_m)^2 + O(\delta_n). \quad (4.7)$$

For the case $\mu \rightarrow a_+$ we get

$$f_m(\mu) = (a_+ - \nu_m)^2 + o(1), \quad m = 1, \dots, N, \quad \text{and} \quad f_m(\mu) = a_-^2 + o(1), \quad m = j, j'. \quad (4.8)$$

Hence the function $f_j - f_{j'}$ (maybe) has the zeros, but the functions $f_j - f_m, m \neq j, j'$ have not zeros in the neighborhood of the point a_+ .

Note that these functions are real outside the small neighborhood of a_+ , otherwise for any complex branches there exists a complex conjugate branch, but the asymptotics (4.7) show that such branches are absent.

We have two cases: (1) let $f_s(\mu), s = j, j'$ be real in some small neighborhood of a_+ . Then the function $f_j - f_{j'}$ has at least one real zero since by Theorem 1.1 the functions $f_j, f_{j'}$ are strongly monotone. Thus $(f_j - f_{j'})^2$ has at least 2 real zeros.

(2) Let $f_s(\mu), s = j, j'$ be complex in some small neighborhood of a_+ . Then they have at least two real branch points. Thus $(f_j - f_{j'})^2$ has at least 2 real zeros.

Hence $(f_j - f_{j'})^2$ has exactly two real zeros, since the number of resonances (in the neighborhood of the point πn) is equal to $N_s(N_s - 1)$.

We determine the sharp asymptotics of resonances. Recall that

$$A = \frac{(-1)^n \Psi(1, z) - I}{-i\varepsilon} = J_1(\mathcal{V} - \mu) - i\widehat{V}_n' + O(\varepsilon), \quad \mathcal{V} = \mathcal{V}_0 \oplus \mathcal{V}_0.$$

The operator $A - a_-$ has the eigenvalue $\xi_0 = \frac{(-1)^n \tau_{n,s} - 1}{i\varepsilon} - a_-$ of multiplicity two, since $\tau_{n,s} = (-1)^n e^{i\varepsilon(a_- + o(1))}$. The operator $J_1(\mathcal{V} - a_+) - a_- = (\mathcal{V}_0 - \nu_j) \oplus (\nu_{j'} - \mathcal{V}_0)$ has two eigenvalue ($= 0$) and other eigenvalues are not zeros. Using the simple transformation (unitary), i.e., changing the lines and columns, $\mu = a_+ + r \in \mathbb{R}$, $r = \tilde{r}_{n,s} \rightarrow 0$, we obtain

$$\det(A - a_- - \xi) = \det \begin{pmatrix} A_1 & A_2 \\ A_3 & A_4 \end{pmatrix} = \det A_4 \det K,$$

$$A_1 = \begin{pmatrix} -r - \xi & ib \\ i\bar{b} & r - \xi \end{pmatrix} + O(\varepsilon), \quad K = A_1 - A_2 A_4^{-1} A_3 = \begin{pmatrix} -r - \xi + a_1 & ib - \xi + a_4 \\ i\bar{b} + a_3 & r + a_2 \end{pmatrix}$$

$$A_4 = \mathcal{V}_{0j} \oplus \mathcal{V}_{0j'} + O(\delta_n), \quad b = -\widehat{v_{n,\alpha}'}, \quad \alpha = (j, j'), \quad A_2, A_3 = O(\delta_n),$$

the function $a_1, a_2, a_3, a_4 = O(\phi)$, $\phi = |\varepsilon| + |\widehat{V}_n'|$ and they analytic with respect to ξ in some small disk. The function $\det K$ has the form

$$\det K = \xi^2 - r^2 + |b|^2 + a_1(r - \xi) + a_2(-r - \xi) - i v a_3 - i a_4 \bar{v} - a_4 a_3 \quad (4.9)$$

Then $0 = \det K = (\xi - \xi_0)^2(1 + O(\xi - \xi_0))$ for $\xi \rightarrow 0$ where $\xi_0 = \frac{((-1)^n \tau_{n,s} - 1)}{i\varepsilon} - a_-$ is the zero of F of multiplicity two. Then $\xi_0 = O(\phi)$ and we have $(r - \gamma)^2 = (|b| - \beta)^2 + O(\phi^2)$ where $\gamma, \beta = O(\phi)$. Then using the estimate $\sqrt{x^2 + y^2} - x \leq y$ for $x, y \geq 0$ we get $r = \pm|b| + O(\phi)$. ■

Proof of Theorem 1.3 We consider N_g^+ , the proof for N_g^- is similar.

(i) Assume that $N_g^+ = \infty$. Then, due to the Lyapunov-Poincarè Theorem and Theorem 1.2, there exists a real sequence $z_k \rightarrow \infty$ as $k \rightarrow \infty$, such that $z_k \in g_{\{j(m), m\}}^{n_k}$ for each $m = 1, \dots, N$. Hence, $\cap_{m=1}^N g_{\{j(m), m\}}^{n_k} \neq \emptyset$. Using asymptotics (1.18) and $k \rightarrow \infty$, we obtain $\nu_1 + \nu_{j(1)} = \dots = \nu_N + \nu_{j(N)}$. Moreover, the estimates $\nu_1 < \dots < \nu_N$ yield $\nu_{j(1)}^0 > \dots > \nu_{j(N)}^0$, i.e. $j(1) = N, j(2) = N - 1, \dots$. Then, $\nu_1 + \nu_N = \nu_2 + \nu_{N-1} = \dots$, which gives a contradiction.
(ii) Let $2a = \nu_1 + \nu_N = \nu_2 + \nu_{N-1} = \dots$. Due to (1.18), $\pi n_k + a \in \cap_{m=1}^N g_{\{N+1-m, m\}}^{n_k}$ as $k \rightarrow \infty$. Then the Lyapunov-Poincarè Theorem yields $\pi n_k + a \notin \sigma(\mathcal{K})$, $k \rightarrow \infty$, i.e. $N_g = \infty$. ■

Proof of Theorem 1.5. i) We need the following result from [K4]: Let $V \in \mathcal{H}$. Define the quasimomentum $k_{j+N} = k_j = \arccos \Delta_j(z) = i \log \eta(\Delta_j(z))$, $j = 1, \dots, N$, see (1.23), (1.22). Then the averaged quasimomentum $k = \frac{1}{2N} \sum_1^{2N} k_j = \frac{1}{N} \sum_1^N k_j$ is analytic in $\mathbb{C}_+$ and $k : \mathbb{C}_+ \rightarrow k(\mathbb{C}_+) = \mathbb{K}(h)$ is a conformal mapping for some $h \in C_{us}$. Furthermore, and there exist branches $k_j, j \in \overline{1, N}$ such that (1.26)-(1.28) hold true.

ii) Let $V, V' \in \mathcal{H}$. We need the following results from [K4]: let for some constants C_0, C_1, C_2 the following asymptotics hold

$$\det(M(z) + M^{-1}(z)) = \exp -i2N \left(z - \frac{C_0}{z} - \frac{C_1}{z^2} - \frac{C_2 + o(1)}{z^3} \right), \quad \text{as } z = iy, \quad y \rightarrow \infty. \quad (4.10)$$

Then

$$k(z) = z - \frac{Q_0}{z} - \frac{Q_1}{z^2} - \frac{Q_2 + o(1)}{z^3}, \quad \text{as } y > r_0|x|, \quad y \rightarrow \infty, \quad \text{for any } r_0 > 0, \quad (4.11)$$

where $C_j = Q_j$, $j = 0, 1, 2$, $Q_2 = I_1^D + I_2^S - \frac{Q_0^2}{2}$. Using these results and asymptotics from Lemma 6.1 we obtain (1.29)-(1.30). ■

5 Example of complex resonances

Let below $N = 2$. Consider the operator $\mathcal{K}_{\nu,\tau} = -iJ_1 \frac{d}{dt} + V_{\nu,\tau}$, $\nu = 1, \frac{1}{2}, \frac{1}{3}, \dots, \tau \in \mathbb{R}$ acting in $L^2(\mathbb{R})^4$, where the real periodic potential $V_{\nu,\tau}$ is given by

$$V_{\nu,\tau} = \begin{pmatrix} 0 & v_{\nu,\tau} \\ v_{\nu,\tau} & 0 \end{pmatrix}, \quad v_{\nu,\tau} = - \begin{pmatrix} a & \tau b_\nu(t) \\ \tau b_\nu(t) & 0 \end{pmatrix}, \quad \frac{a}{2\pi} \in \mathbb{R}_+ \setminus \mathbb{N}, \quad b_\nu \in C(\mathbb{T}). \quad (5.1)$$

We need another representation of $\mathcal{K}_{\nu,\tau}$. Recall that $\mathcal{U} = \frac{1}{\sqrt{2}}(J_1 + iJ) = \mathcal{U}^*$ and identities (2.22) give $\mathcal{K}_{\nu,\tau}^- = \mathcal{U} \mathcal{K}_{\nu,\tau} \mathcal{U} = J \frac{d}{dt} - V_{\nu,\tau}$. Using the unitary transformation $y = (y_1, y_2, y_3, y_4)^\top \rightarrow \mathcal{U}y = (y_1, y_3, y_2, y_4)^\top$ in $L^2(\mathbb{R})^4$ we define the new operator $P_{\nu,\tau} = \mathcal{U} \mathcal{K}_{\nu,\tau}^- \mathcal{U}^* = \mathbf{j} \frac{d}{dt} + W_{\nu,\tau}$, where

$$W_{\nu,\tau} = -\mathcal{U} V_{\nu,\tau} \mathcal{U}^* = \begin{pmatrix} a\mathbf{j}_2 & \tau b_\nu \mathbf{j}_2 \\ \tau b_\nu \mathbf{j}_2 & 0 \end{pmatrix}, \quad \mathbf{j}_2 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \mathbf{j} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \quad (5.2)$$

We rewrite $W_{\nu,\tau}$ in the form $W_{\tau,\nu} = W^0 + \tau b_\nu J_2$, where $W^0 = a \begin{pmatrix} \mathbf{j}_2 & 0 \\ 0 & 0 \end{pmatrix}$. If $\tau = 0$, then we have the unperturbed operator $P^0 = \mathbf{j} \frac{d}{dt} + W^0$ with a constant potential W^0 .

1. The 2×2 Dirac operator. We consider the simple example of the 2×2 Dirac operator $P_1^0 = \mathbf{j} \frac{d}{dt} + a\mathbf{j}_2$, $a > 0$ acting in $L^2(\mathbb{R}) \oplus L^2(\mathbb{R})$. The spectrum of P_1^0 is purely absolutely continuous and consists of two intervals $(-\infty, -a), (a, \infty)$ separated by the gap $(-a, a)$. In this case only one gap is open and other gaps are closed. The solution of the system $\mathbf{j}y' + a\mathbf{j}_2 y = zy$ has the form $y = e^{\pm ikt} y_0$, for some constant vector $y_0 \in \mathbb{C}^2$ and k satisfies

$$\det(ik\mathbf{j} + a\mathbf{j}_2 - zI_2) = -k^2 - a^2 + z^2, \quad k = k(z) = \sqrt{z^2 - a^2},$$

where the quasimomentum $k : \mathbb{C} \setminus [-a, a] \rightarrow \mathbb{C} \setminus [ia, -ia]$ is the conformal mapping with asymptotics $k(z) = z + o(1)$ as $|z| \rightarrow \infty$. Note that

$$A = \mathbf{j}(z - a\mathbf{j}_2), \quad A^2 = (\mathbf{j}(z - a\mathbf{j}_2))^2 = -z^2 + a^2 - za\mathbf{j}\mathbf{j}_2 - za\mathbf{j}\mathbf{j}_2 = a^2 - z^2 = -k^2. \quad (5.3)$$

Then the fundamental solution of the equation $\mathbf{j}\psi_1^{0'} + a\mathbf{j}_2\psi_1^0 = z\psi_1^0$, $\psi_1^0(0, z) = I_2$ is given by

$$\psi_1^0(t, z) = e^{-tA} = \sum_{n \geq 0} \frac{(-tA)^n}{n!} = \sum_{n \geq 0} \left(\frac{(tA)^{2n}}{(2n)!} - \frac{(tA)^{2n+1}}{(2n+1)!} \right)$$

$$= \sum_{n \geq 0} \left(\frac{(-1)^n (tk)^{2n}}{(2n)!} - \frac{A}{k} \frac{(-1)^n (tk)^{2n+1}}{(2n+1)!} \right) = \cos tk - \frac{A}{k} \sin tk. \quad (5.4)$$

Thus the Lyapunov function has the form

$$\Delta_1^0(z) = \frac{\text{Tr } e^{-A}}{2} = \cos k(z), \quad k = k(z) = \sqrt{z^2 - a^2}, \quad z \in \mathbb{C} \setminus [-a, a]. \quad (5.5)$$

2. The 4×4 unperturbed operator. Consider the 4×4 operator $P^0 = \mathbf{j} \frac{d}{dt} + W^0$ in $L^2(\mathbb{R})^2 \oplus L^2(\mathbb{R})^2$, where $W^0 = \begin{pmatrix} a\mathbf{j}_2 & 0 \\ 0 & 0 \end{pmatrix}$ is the 4×4 matrix. We rewrite this operator in the form $P^0 = P_1^0 \oplus P_2^0$, where the Dirac operators $P_1^0 = \mathbf{j} \frac{d}{dt} + a\mathbf{j}_2$ and $P_2^0 = \mathbf{j} \frac{d}{dt}$ act in $L^2(\mathbb{R}) \oplus L^2(\mathbb{R})$. The corresponding equation $\mathbf{j} \frac{d}{dt} \psi^0 + W^0 \psi^0 = z \psi^0$ has the fundamental solution $\psi^0(t)$ given by

$$\psi^0(t) = \psi_1^0(t) \oplus \psi_2^0(t), \quad \psi_1^0 = e^{-At}, \quad \psi_2^0 = e^{-\mathbf{j}zt}, \quad A = \mathbf{j}(z - a\mathbf{j}_2), \quad t \geq 0. \quad (5.6)$$

Thus using (5.4)-(5.6), (1.11) we obtain that $D^0(\tau, \cdot) = \det(\psi^0(1) - \tau I_4)$ satisfies

$$D^0(\tau, \cdot) = (\tau^2 - 2\Delta_1^0 \tau + 1)(\tau^2 - 2\Delta_2^0 \tau + 1), \quad \Delta_m^0 = (T_1^0 - (-1)^m \sqrt{\rho^0})/2,$$

$$T_m^0(z) = \text{Tr}(e^{-mA} + e^{-\mathbf{j}zm}) = 2(\cos mk(z) + \cos mz), \quad \rho^0(z) = (\cos k(z) - \cos z)^2, \quad (5.7)$$

for any $z \in \mathbb{C}, m = 1, 2$. We will determine the zeros of $\rho^0(z)$. We have $0 = \cos k - \cos z = 2 \sin \frac{k+z}{2} \sin \frac{k-z}{2}$, $k = k(z)$. Then we obtain $k \pm z = 2\pi n, n \in \mathbb{Z} \setminus \{0\}$, which gives zeros r_n^0 (each zero has multiplicity 2) of ρ^0 by

$$r_n^0 = \pi n + \frac{a^2}{4\pi n}, \quad k(r_n^0) = \begin{cases} \pi n - \frac{a^2}{4\pi n} & \text{if } |n| > \frac{a}{2\pi} \\ -\pi n + \frac{a^2}{4\pi n} & \text{if } |n| < \frac{a}{2\pi} \end{cases}, \quad n \in \mathbb{Z} \setminus \{0\}. \quad (5.8)$$

We determine the periodic spectrum for the equation $\mathbf{j}y' + W^0 y = zy$. Using (5.6), (5.7) we have $\det(\psi^0(1, z) \mp I_4) = 4(\cos k(z) \mp 1)(\cos z \mp 1)$, which yields the periodic and anti-periodic spectrum multiplicity 2

$$z_{n,1}^{0,\pm} = \pi n, n \in \mathbb{Z}, \quad \text{and} \quad z_{n,2}^{0,\pm} = \pi n \sqrt{1 + \frac{a^2}{\pi^2 n^2}}, \quad n \in \mathbb{Z} \setminus \{0\}, \quad (5.9)$$

and $z_{0,2}^{0,\pm} = \pm a$ has multiplicity one. Note that the zeros $z_{n,p}^0 \neq r_m^0$ for all $n, m \in \mathbb{Z}, p = 1, 2$. In this case there are no gaps in the spectrum.

3. The perturbed case. Consider the 4×4 operator $P_{\tau,\nu} = \mathbf{j} \frac{d}{dt} + W_{\tau,\nu}$ in $L^2(\mathbb{R})^2 \oplus L^2(\mathbb{R})^2$, where the 4×4 potential $W_{\tau,\nu} = W^0 + \tau b_\nu \mathbf{j}_2 J_2$ satisfies (5.1), (1.20). We show that there exist the non-degenerated resonance gaps for some $W_{\tau,\nu}$. In this example some resonances are real and some are complex. The fundamental solution $\psi^{\tau,\nu}(t, z)$ of the Eq. $\mathbf{j}\psi' + W_{\tau,\nu}\psi = z\psi$ satisfies the integral equation

$$\psi^{\tau,\nu}(t, z) = \psi^0(t, z) + \tau \int_0^t \psi^0(t-s, z) \mathbf{j} b_\nu(s) \mathbf{j}_2 J_2 \psi^{\tau,\nu}(s, z) ds. \quad (5.10)$$

Then $\psi^{\tau,\nu}(t, z)$ has asymptotics

$$\psi^{\tau,\nu}(m, z) = \psi^0(m, z) + \tau\psi^1(m, z, \nu) + \tau^2\psi^2(m, z, \nu) + O(\tau^3 e^{m|\operatorname{Im} z|}) \quad \text{as } \tau \rightarrow 0, \quad (5.11)$$

uniformly in $\nu = 1, \frac{1}{2}, \dots, m = 1, 2, z \in \mathbb{C}$, where

$$\psi^1(m, z, \nu) = \int_0^m \psi^0(m-t, z) b_\nu(t) \mathbf{j}_1 J_2 \psi_0(t, z) dt, \quad (5.12)$$

$$\psi^2(m, z, \nu) = \int_0^m \psi^0(m-t, z) b_\nu(t) \mathbf{j}_1 J_2 dt \int_0^t \psi^0(t-s, z) b_\nu(s) \mathbf{j}_1 J_2 \psi_0(s, z) ds. \quad (5.13)$$

The identity $\operatorname{Tr} \psi^0(t, z) \mathbf{j}_1 J_2 = 0$, $(t, z) \in \mathbb{R} \times \mathbb{C}$ yields $\operatorname{Tr} \psi^1(t, z, \nu) = 0$. Thus we obtain

$$T_m^{\tau,\nu}(z) \equiv \operatorname{Tr} \psi^{\tau,\nu}(m, z) = T_m^0(z) + \tau^2 T_{m2}(z, \nu) + O(\tau^3 e^{m|\operatorname{Im} z|}), \quad m = 1, 2, \quad (5.14)$$

$$T_{m2}(z, \nu) = \int_0^m b_\nu(t) dt \int_0^t b_\nu(s) F_m(t, s, z) ds, \quad F_m(t, s, z) = \operatorname{Tr} \mathbf{j}_1 J_2 \psi^0(y, z) \mathbf{j}_1 J_2 \psi_0(\zeta, z), \quad (5.15)$$

uniformly in $\nu = 1, \frac{1}{2}, \dots, |\tau| < 1, z \in \mathbb{C}$, where $y = t - s, \zeta = m - y$.

Lemma 5.1. *Let $V_{\tau,\nu}$ satisfy (1.19), (1.20). Then the following asymptotics hold true*

$$T_{12}(z, \nu) = \frac{T_1^0(z)}{2} + o(e^{|\operatorname{Im} z|}), \quad T_{22}(z, \nu) = T_2^0(z) + 4\phi(z) + o(e^{2|\operatorname{Im} z|}), \quad (5.16)$$

$$T_1^{\tau,\nu}(z) = T_1^0(z) \left(1 + \frac{\tau^2}{2}\right) + o(\tau^2 e^{|\operatorname{Im} z|}), \quad (5.17)$$

$$T_2^{\tau,\nu}(z) = (1 + \tau^2) T_2^0(z) + \tau^2 4\phi(z) + o(\tau^2 e^{2|\operatorname{Im} z|}), \quad (5.18)$$

$$\rho^{\tau,\nu}(z) = \frac{T_2^{\tau,\nu}(z) + 4}{2} - \frac{T_1^{\tau,\nu}(z)^2}{4} = (1 + \tau^2) \rho^0(z) + \tau^2 2(\phi(z) - 1) + o(\tau^2 e^{2|\operatorname{Im} z|}), \quad (5.19)$$

$$\det(\psi^{\tau,\nu}(1, z) \mp I_4) = (1 + \tau^2) D^0(\pm 1, z) + \tau^2 \left(T_1^0 - 1 - \frac{\phi(z) - 1}{2} + o(e^{2|\operatorname{Im} z|}) \right), \quad (5.20)$$

as $\nu \rightarrow 0$, uniformly on $|\tau| < 1, z \in \mathbb{C}$, and where $\phi(z) = \cos k(z) \cos z + \frac{z}{k(z)} \sin z \sin k(z)$.

Proof. Using $b_\nu \rightarrow \delta_{per} = \sum \delta(t - \frac{1}{2} - n)$ and $F_j(\cdot, \cdot) \in C(\mathbb{R}^2), j = 1, 2$, we obtain

$$\int_0^1 dt \int_0^t b_\nu(t) b_\nu(s) F_1(t, s) ds = \frac{1}{2} F_1\left(\frac{1}{2}, \frac{1}{2}\right) + o(e^{|\operatorname{Im} z|}), \quad (5.21)$$

$$\int_0^2 dt \int_0^t b_\nu(t) b_\nu(s) F_2(t, s) ds = \frac{1}{2} F_2\left(\frac{1}{2}, \frac{1}{2}\right) + \frac{1}{2} F_2\left(\frac{3}{2}, \frac{3}{2}\right) + F_2\left(\frac{3}{2}, \frac{1}{2}\right) + o(e^{2|\operatorname{Im} z|}) \quad (5.22)$$

as $\nu \rightarrow 0$. Thus, if $m = 1$, then (5.21) gives

$$T_{12}(z, \nu) = \frac{1}{2} \operatorname{Tr} \mathbf{j}_1 J_2 \psi^0(0, z) \mathbf{j}_1 J_2 \psi_0(1, z) + o(e^{|\operatorname{Im} z|}) = \frac{T_1^0(z)}{2} + o(e^{|\operatorname{Im} z|}),$$

If $m = 2$, then $F_2(\frac{1}{2}, \frac{1}{2}) = F_2(\frac{3}{2}, \frac{3}{2}) = \text{Tr } \psi_0(2, z)$ and the identity $J_2\psi^0 = \begin{pmatrix} 0 & \psi_2^0 \\ \psi_1^0 & 0 \end{pmatrix}$ yields

$$\begin{aligned} F_2\left(\frac{3}{2}, \frac{1}{2}\right) &= \text{Tr } \mathbf{j}_1 \begin{pmatrix} 0 & \psi_2^0(y) \\ \psi_1^0(y) & 0 \end{pmatrix} \mathbf{j}_1 \begin{pmatrix} 0 & \psi_2^0(\zeta) \\ \psi_1^0(\zeta) & 0 \end{pmatrix} = 2 \text{Tr } \mathbf{j}_1 \psi_2^0(1, z) \mathbf{j}_1 \psi_1^0(1, z) \\ &= 2 \text{Tr } \mathbf{j}_1 (\cos z - \mathbf{j} \sin z) \mathbf{j}_1 (\cos k - \frac{A}{k} \sin k) = 2 \text{Tr} (\cos z + \mathbf{j} \sin z) (\cos k - \frac{A}{k} \sin k) \\ &= 4 \cos z \cos k - 2 \text{Tr } \mathbf{j} \frac{A}{2k} \sin z \sin k = 4(\cos z \cos k + \frac{z}{k} \sin z \sin k) = 4\phi. \end{aligned}$$

Then (5.22) gives

$$T_{22}(z, \nu) = \text{Tr } \psi_0(2, z) + 4\phi(z) = T_2^0(z) + 4\phi(z) + o(e^{2|\text{Im } z|}) \quad \text{as } \nu \rightarrow 0.$$

Substituting (5.16) into (5.14) we obtain (5.17), (5.18). The asymptotics (5.17), (5.18) imply

$$\rho^{\tau, \nu}(z) = \frac{T_2^{\tau, \nu}(z) + 4}{2} - \frac{T_1^{\tau, \nu}(z)^2}{4} = (1 + \tau^2)\rho^0(z) + \tau^2 2(\phi(z) - 1) + o(\tau^2 e^{2|\text{Im } z|}),$$

and $D^{\tau, \nu}(\pm 1, \cdot) = \det(\psi^{\tau, \nu}(1, \cdot) \mp I_4) = (T_1^{\tau, \nu} - 1)^2 - \rho^{\tau, \nu}$ satisfies

$$\begin{aligned} D^{\tau, \nu}(1, z) &= \left((1 + \frac{\tau^2}{2}) T_1^0(z) - 1 \right)^2 - (1 + \tau^2)\rho^0(z) - \frac{\tau^2}{2}(\phi(z) - 1) + o(\tau^2 e^{|\text{Im } z|}) \\ &= D^0(1, z) + \tau^2 \left(T_1^0(z)(T_1^0(z) - 1) - \rho^0(z) - \frac{(\phi(z) - 1)}{2} + o(1) \right) \end{aligned}$$

as $\nu \rightarrow 0$, which gives (5.20). The proof for $D^{\tau, \nu}(-1, \cdot)$ is similar. ■

Proof of Proposition 1.4. We have the simple asymptotics

$$k(z) = z - \frac{a^2}{2z} + O(z^{-3}), \quad \cos k(z) = \cos z + \frac{a^2}{2z} \sin z + O(z^{-2} e^{|\text{Im } z|}), \quad (5.23)$$

$$\rho^0(z) = (\cos k(z) - \cos z)^2 = \left(\frac{a^2 \sin z}{2z} \right)^2 + O(z^{-3} e^{2|\text{Im } z|}) \quad (5.24)$$

as $|z| \rightarrow \infty$. We take $\varkappa > 0$ such that the disks $B_n = \{|z - r_n^0| < \varkappa\}$, $n \in \mathbb{Z}$ are not overlapping. Define a constants $\eta_n = \min_{|z - r_n^0| = \varkappa} |\rho^0(z)| > 0$. Thus using (5.20) we get

$$|\rho^{\tau, \nu}(z) - \rho^0(z)| = \tau^2 O(1) = \frac{\tau^2}{\eta_n} \rho^0(z) O(1), \quad |z - r_n^0| = \varkappa \quad \text{as } \tau \rightarrow 0, \quad (5.25)$$

for each n . We also obtain for $|z| = \pi n_0 + 1$ the following estimates

$$|\rho^{\tau, \nu}(z) - \rho^0(z)| = \tau^2 e^{2|\text{Im } z|} O(1) = \tau^2 |4 \sin z|^2 O(1) = \tau^2 |z|^2 |\rho^0(z)| O(1), \quad n_0 \rightarrow \infty. \quad (5.26)$$

Thus we take large n_0 and sufficiently small τ such that $|\rho(z) - \rho^0(z)| \leq \frac{1}{2} |\rho^0(z)|$ on all contours $|z| = \pi(n_0 + 1)$ and $|z - r_n^0| = \varkappa$, $|n| \leq n_0$. Then by the Rouché theorem, ρ has as

many roots, counted with multiplicity, as ρ^0 in the disks $\{|z| < \pi n_0 + 1\}, \{|z - r_n^0| < \varkappa\}$. Since ρ^0 has exactly one double root at $r_n^0, n \neq 0$, and since n_0 can be chosen arbitrarily large, we deduce that in each disk $\{|z - r_n^0| < \varkappa\}, 1 \leq |n| \leq n_0$ there exist two zeros $r_{n,\nu}^\pm(\tau)$ of $\rho^{\tau,\nu}$ for sufficiently small τ, ν .

Consider the zeros $r_{n,\nu}^\pm(\tau)$ of $\rho^{\tau,\nu}$ in the disk $\{|z - r_n^0| < \varkappa\}$ for fixed $n, 1 \leq n \leq n_0$. The proof for the case $n < 0$ is similar. Recall $\phi(z) = \cos k(z) \cos z + \frac{z}{k(z)} \sin z \sin k(z)$. Consider $\phi(z) - 1$ at the point $r_n^0 = \pi n + x_n, n \geq 1$, where $x_n = \frac{a^2}{4\pi n}$. Using (5.8) we obtain

$$\phi(r_n^0) - 1 = \cos^2 x_n - 1 + \frac{r_n^0}{k(r_n^0)} \sin^2 x_n = \left(\frac{r_n^0}{k(r_n^0)} - 1 \right) \sin^2 x_n > 0, \quad \text{if } 1 \leq n < \frac{a}{2\pi}, \quad (5.27)$$

$$\phi(r_n^0) - 1 = \cos^2 x_n - 1 - \frac{r_n^0}{k(r_n^0)} \sin^2 x_n = -\left(1 + \frac{r_n^0}{k(r_n^0)} \right) \sin^2 x_n < 0, \quad \text{if } n > \frac{a}{2\pi}. \quad (5.28)$$

We rewrite the function $\rho^{\tau,\nu}$ in the disk $\{|z - r_n^0| < \varkappa\}$ in the form

$$R(z, \tau) \equiv \frac{\rho^{\tau,\nu}(z)}{(1 + \tau^2)^2} = (z - r_n^0)^2 f(z) + \tau^2 \phi_1(z, \tau), \quad f(z) = \frac{\rho^0(z)}{(z - r_n^0)^2}, \quad z \in B_n, \quad (5.29)$$

for sufficiently small fixed ν, τ . The functions $R(z, \tau), f(z), \phi_1(z, \tau)$ are analytic in $(z, \tau) \in B_n \times \{|\tau| < \varepsilon\}$ for some small $\varkappa, \varepsilon > 0$ and satisfy

$$f(r_n^0) > 0, \quad \phi_1(r_n^0, 0) = 2(\phi(r_n^0) - 1) + o(1) \quad \text{as } \nu \rightarrow 0. \quad (5.30)$$

Applying the Implicit Function Theorem to $R(z, \tau) = 0$ and using (5.27)-(5.30) we obtain a unique solution $r_{n,\nu}^\pm(\tau)$ of the equation $\Phi(r_{n,\nu}^\pm(\tau), \tau) = 0, \tau \in (-\tau_0, \tau_0), r_{n,\nu}^\pm(0) = r_n^0$, for some $\tau_0 > 0$ and here $r_{n,\nu}^\pm(\tau)$ is an analytic function in $\{|\tau| < \tau_0\}$ and satisfies (1.21) with $R_n = -2(\phi(r_n^0) - 1)/f(r_n^0)$. The proof of the statement ii) is similar. ■

6 Appendix

Lemma 6.1. *Let $V, V' \in \mathcal{H}$ and let $r > 0$. Then for $y \geq r|x|, y \rightarrow \infty$ following asymptotics hold:*

$$\text{Tr } L_1(z) = \|V\|^2 \sin z, \quad (6.1)$$

$$\text{Tr } L_2(z) = \left(i\mathcal{H}_1 - \frac{\text{Tr } \mathcal{V}^2}{2} + i \frac{\|V'\|^2 + o(1)}{2z} \right) \cos z, \quad (6.2)$$

$$\text{Tr } \mathcal{V} L_2(z) = \cos z \text{Tr} \left(G_1 + G - \frac{\mathcal{V}^3}{2} \right) + o(e^{|\text{Im } z|}), \quad (6.3)$$

$$\text{Tr } L_3(z) = i \cos z \text{Tr} \left(\int_0^1 V_t^4 dt + G_1 + G - \frac{\mathcal{V}^3}{6} \right) + o(e^{|\text{Im } z|}), \quad (6.4)$$

$$G_1 = \mathcal{V} \int_0^1 J_1 V_t' V_t dt, \quad G = \mathcal{V} \int_0^1 J_1 u_t' u_t dt, \quad u_t = \int_0^t V_s^2 ds,$$

$$\det L(z) = 2^{-2N} \exp -i \left(2Nz - \frac{\mathcal{H}_0}{2z} - \frac{\mathcal{H}_1}{(2z)^2} - \frac{\mathcal{H}_2 + o(1)}{(2z)^3} \right), \quad (6.5)$$

where $\mathcal{H}_0 = \text{Tr} \int_0^1 V_t^2 dt$, $\mathcal{H}_1 = \text{Tr} \int_0^1 -iJ_1 V_t' V_t dt$, $\mathcal{H}_2 = \text{Tr} \int_0^1 (V_t'^2 + V_t^4) dt$.

Proof. Identity (3.15) implies (6.1). We will show (6.2). Asymptotics (3.12) yields

$$L_2 = -i \sin z \int_0^1 J_1 V_t' V_t dt + i \frac{e_1 J_1}{2} (\mathcal{V} \widehat{V'} + \widehat{V'} \mathcal{V}) + L_{21} + L_{22}, \quad (6.6)$$

$$L_{21} = -\frac{1}{2} \int_0^1 (e_1 u' u + e_{-1} u u') dt, \quad L_{22} = \frac{1}{2} \int_0^1 (e_1 f' f + e_{-1} f f') dt, \quad f_t = \int_0^t V_s' e_{2s} ds$$

as $|z| \rightarrow \infty, y \geq r|x|$. We determine $\text{Tr} L_2$. Using (2.12) we have $\text{Tr} e_1 J_1 (\mathcal{V} \widehat{V'} + \widehat{V'} \mathcal{V}) = 0$ and then

$$\text{Tr} L_2 = -i \sin z \mathcal{H}_1 + \text{Tr}(L_{21} + L_{22}). \quad (6.7)$$

Using (2.15) we get

$$\text{Tr} L_{21} = -\frac{\text{Tr}}{2} \int_0^1 (e_1 u' u + e_{-1} u u') dt = -\frac{\cos z}{2} \text{Tr} \int_0^1 (u' u + u u') dt = -\frac{\cos z}{2} \text{Tr} \mathcal{V}^2. \quad (6.8)$$

Due to (2.15), (2.2) we obtain

$$\text{Tr} L_{22} = \frac{\text{Tr}}{2} \int_0^1 (e_1 f' f + e_{-1} f f') dt = \text{Tr} \int_0^1 \int_0^t e^{izJ_1(1-2t+2s)} V_t' V_s' dt ds \quad (6.9)$$

and using

$$\int_0^1 \int_s^t e^{i2z(t-s)} f_t r_s dt ds = \int_0^1 \frac{i f_t' r_t dt}{2z} + \frac{o(1)}{z}, \quad \int_0^1 \int_s^t e^{-i2z(t-s)} f_t r_s dt ds = \frac{o(e^{|\text{Im} z|})}{z}, \quad (6.10)$$

for $f, h \in L^2(0, 1)$ and $\text{Tr} J_1 (V')^2 = 0$, we have $\text{Tr} L_{22} = \frac{i \cos z}{2z} (\|V'\|^2 + o(1))$, which give (6.2). We will determine $\text{Tr} \mathcal{V} L_2$. Using (6.6) we have

$$\text{Tr} \mathcal{V} L_2 = \text{Tr} \mathcal{V} \left(-i \sin z \int_0^1 J_1 V' V dt + L_{21} \right) + o(e^{|\text{Im} z|}), \quad (6.11)$$

since $\text{Tr} \mathcal{V} e_1 J_1 (\mathcal{V} \widehat{V'} + \widehat{V'} \mathcal{V}) = 0$ and $L_{22}(z) = o(e^{|\text{Im} z|})$. Due to $e^{izJ_1} = \cos z + iJ_1 \sin z$ and $\int_0^1 (u' u + u u') dt = \mathcal{V}^2$, $\text{Tr} \mathcal{V}^3 = 0$ we get

$$\text{Tr} \mathcal{V} L_{21} = -\frac{\cos z}{2} \text{Tr} \mathcal{V}^3 - i \frac{\sin z}{2} \text{Tr} \mathcal{V} J_1 \int_0^1 (u' u - u u') dt = -\frac{\cos z}{2} \text{Tr} \mathcal{V}^3 - i \sin z G$$

and together with (6.11) we obtain (6.3), since $\sin z = i \cos + O(e^{-|\text{Im} z|})$.

We will determine $\text{Tr } L_3$. Using (3.13) we rewrite $\text{Tr } L_3$ in the form

$$\text{Tr } L_3 = \text{Tr } \varkappa_3 = \text{Tr}(KV^2 + K^2V + KVK + K^3)\psi_0 = \sum_1^4 A_k,$$

$$A_1 = \text{Tr } KV^2\psi_0, \quad A_2 = \text{Tr } K^2V\psi_0, \quad A_3 = \text{Tr } KVK\psi_0, \quad A_4 = \text{Tr } K^3\psi_0, \quad (6.12)$$

and recall $(Kf)(t) = \int_0^t e_{t-s}W_s f(s)ds$, $W = -iJ_1V^2 - V'$. Using (2.12), (2.14), (2.15) and $e_1 = \cos z + iJ_1 \sin z$ we have

$$\begin{aligned} A_1 &= \text{Tr} \int_0^1 e_1 W_t V_t^2 dt = \text{Tr} \int_0^1 e_1 (-iJ_1 V_t^2 - V_t') V_t^2 dt = \text{Tr} \int_0^1 -iJ_1 e_1 V_t^4 dt = \\ &= \text{Tr} \int_0^1 -iJ_1 (\cos z + iJ_1 \sin z) V_t^4 dt = \sin z \text{Tr} \int_0^1 V_t^4 dt. \end{aligned}$$

The similar arguments give

$$\begin{aligned} A_2 &= \text{Tr} \int_0^1 \int_0^t e_{1-t+s} W_t e_{t-s} W_s V_s dt ds = \text{Tr} \int_0^1 \int_0^t e_{1-t+s} (iJ_1 V_t^2 + V_t') e_{t-s} (iJ_1 V_s^2 + V_s') V_s dt ds \\ &= \text{Tr} \int_0^1 \int_0^t iJ_1 (e_1 V_t^2 V_s' V_s - e_{1-2t+2s} V_t' V_s^3) dt ds \end{aligned} \quad (6.13)$$

and

$$\begin{aligned} A_3 &= \text{Tr} \int_0^1 \int_0^t e_{1-t+s} W_t V_t e_{t-s} W_s dt ds = \text{Tr} \int_0^1 \int_0^t e_{1-t+s} (iJ_1 V_t^2 + V_t') V_t e_{t-s} (iJ_1 V_s^2 + V_s') dt ds \\ &= \text{Tr} \int_0^1 \int_0^t iJ_1 (e_1 V_t' V_t V_s^2 + e_{1-2t+2s} V_t^3 V_s') dt ds. \end{aligned} \quad (6.14)$$

Summing (6.13), (6.14) we get $A_2 + A_3 = F_0 + F_1$, where

$$F_0 = \text{Tr} \int_0^1 \int_0^t iJ_1 e_1 (V_t^2 V_s' V_s + V_t' V_t V_s^2) dt ds = \text{Tr} iJ_1 e_1 \mathcal{V} \int_0^1 V_t' V_t dt = i \cos z \text{Tr} J_1 \mathcal{V} \int_0^1 V_t' V_t dt, \quad (6.15)$$

since $iJ_1 e_1 = iJ_1 \cos z - \sin z$ and $\text{Tr} \mathcal{V} \int_0^1 V_t' V_t dt = 0$. We will show

$$F_1 = \text{Tr} \int_0^1 \int_0^1 iJ_1 e_{1-2t+2s} (-V_t' V_s^3 + V_t^3 V_s') dt ds = o(e^{|\text{Im } z|}). \quad (6.16)$$

We use the standard arguments. If $V, V', V'' \in \mathcal{H}$, then integration by parts gives (6.16). If $V, V' \in \mathcal{H}$, then there exists $P_h, P_h', P_h'' \in \mathcal{H}$ such that $\|V - P_h\| + \|V' - P_h'\| = h$ for some small $h \geq 0$. Then $F_1 = o(e^{|\text{Im } z|})(1 + O(h))$, which yields (6.16), since h is arbitrary small.

The similar arguments give

$$A_4 = \text{Tr} \int_0^1 \int_0^t \int_0^s e_{1-t+p} W_t e_{t-s} W_s e_{s-p} W_p dt ds dp = F_2 + F_3,$$

where

$$F_2 = \text{Tr} \int_0^1 \int_0^t \int_0^p (iJ_1) e_1 V_t^2 V_s^2 V_p^2 dt ds dp,$$

$$F_3 = \text{Tr} \int_0^1 \int_0^t \int_0^p (iJ_1) \left(e_{1+p-s} V_t^2 V_s' V_p' + e_{1-2t+2p} V_t' V_s^2 V_p' + e_{1-2t+2s} V_t' V_s' V_p^2 \right) dt ds dp$$

Using $u_s = \int_0^s V_t^2 dt$ and $\int_s^1 V_t^2 dt = \mathcal{V} - u_s$ we obtain

$$F_2 = \text{Tr}(iJ_1 e_1) \int_0^1 \int_s^1 u_t' u_s' u_s dt ds = \text{Tr}(iJ_1 e_1) \int_0^1 (\mathcal{V} - u_s) u_s' u_s ds = i \cos z \text{Tr} G - \frac{\sin z}{6} \text{Tr} \mathcal{V}^3, \quad (6.17)$$

since

$$\text{Tr}(iJ_1 e_1) \int_0^1 u_s u_s' u_s ds = \text{Tr} \frac{iJ_1 e_1}{3} \int_0^1 (u_s' u_s^2 + u_s u_s' u_s + u_s^2 u_s') ds = \text{Tr} \frac{iJ_1 e_1}{3} \mathcal{V}^3 = -\frac{\sin z}{3} \text{Tr} \mathcal{V}^3$$

and

$$\begin{aligned} \text{Tr}(iJ_1 e_1) \mathcal{V} \int_0^1 u_s' u_s ds &= i \cos z \text{Tr} J_1 \mathcal{V} \int_0^1 u_s' u_s ds - \sin z \text{Tr} \mathcal{V} \int_0^1 u_s' u_s ds \\ &= i \cos z \text{Tr} G - \frac{\sin z}{2} \text{Tr} \mathcal{V} \int_0^1 (u_s' u_s + u_s u_s') ds = i \cos z \text{Tr} G - \frac{\sin z}{2} \text{Tr} \mathcal{V}^3, \end{aligned}$$

where we used: $\text{Tr} J_1 \mathcal{V}^3 = 0$ and $\text{Tr} ABC = \text{Tr} ACB$ for real self-adjoint matrix and real representations (2.23), (2.26) of V . Using standard arguments (see the proof of (6.16)) we have $F_3 = o(e^{|\text{Im} z|})$. Summing $A_1, \dots, A_4$ we have (6.4).

We will determine (6.5). Asymptotics (3.12) yields

$$\frac{L}{\cos z} = I_{2N} + S, \quad S = i\varepsilon \mathcal{V} + \frac{\varepsilon^2 L_2}{\cos z} + \frac{\varepsilon^3 L_3}{\cos z} + O(\varepsilon^4), \quad S = O(\varepsilon) \quad (6.18)$$

as $\text{Im} z \rightarrow \infty$. In order to use the identity

$$\det(I + S) = e^\Phi, \quad \Phi = \text{Tr} S - \text{Tr} \frac{S^2}{2} + \text{Tr} \frac{S^3}{3} + O(\varepsilon^4), \quad |S| = O(\varepsilon),$$

we need the traces of $S^m, m = 1, 2, 3$. Due to (6.18), we get $\text{Tr} \frac{S^3}{3} = -i\varepsilon^3 \text{Tr} \frac{\mathcal{V}^3}{3} + O(\varepsilon^4)$. Using (6.18), (6.1)-(6.4) we get

$$-\text{Tr} \frac{S^2}{2} = \text{Tr} \left(\varepsilon^2 \frac{\mathcal{V}^2}{2} - i\varepsilon^3 \frac{\mathcal{V} L_2}{\cos z} + o(\varepsilon^3) \right) = \text{Tr} \left(\varepsilon^2 \frac{\mathcal{V}^2}{2} - i\varepsilon^3 \left(G_1 + G - \frac{\mathcal{V}^3}{2} + o(\varepsilon^3) \right) \right), \quad (6.19)$$

$$\text{Tr} S = i\varepsilon \mathcal{H}_0 + \varepsilon^2 \text{Tr} \left(i\mathcal{H}_1 - \varepsilon^2 \frac{\mathcal{V}^2}{2} \right) + i\varepsilon^3 \left(\mathcal{H}_2 + G_1 + G - \frac{\mathcal{V}^3}{6} + o(1) \right), \quad (6.20)$$

and summing (??)-(6.20) we get $\Phi = i\varepsilon \mathcal{H}_0 + i\varepsilon^2 \mathcal{H}_1 + i\varepsilon^3 \mathcal{H}_2 + o(\varepsilon^3)$, which yields (6.5). ■

References

[BBK] Badanin, A.; Brüning, J.; Korotyaev, E. The Lyapunov function for Schrödinger operators with a periodic 2×2 matrix potential. J. Funct. Anal. 234 (2006), 106–126

- [CK] Chelkak, D.; Korotyaev, E. Spectral estimates for Schrödinger operators with periodic matrix potentials on the real line. *Int. Math. Res. Not.* 2006, Art. ID 60314, 41 pp.
- [CG] Clark, S.; Gesztesy, F. Weyl-Titchmarsh M -function asymptotics, local uniqueness results, trace formulas, and Borg-type theorems for Dirac operators. *Trans. Amer. Math. Soc.* 354 (2002), no. 9, 3475–3534
- [CHGL] Clark S.; Holden H.; Gesztesy, F.; Levitan, B. Borg-type theorem for matrix-valued Schrödinger and Dirac operators, *J. Diff. Eqs.* 167(2000), 181-210
- [DS] Dunford, N. and Schwartz, J.: *Linear Operators Part II: Spectral Theory*, Interscience, New York, 1988
- [Fo] Forster, O. *Lectures on Riemann surfaces*. Graduate Texts in Mathematics, 81. Springer-Verlag, New York, 1991
- [GL] Gel'fand I.; Lidskii, V. On the structure of the regions of stability of linear canonical systems of differential equations with periodic coefficients. (Russian) *Uspehi Mat. Nauk (N.S.)* 10 (1955), no. 1(63), 3–40
- [Ge] Gel'fand, I. Expansion in characteristic functions of an equation with periodic coefficients. (Russian) *Doklady Akad. Nauk SSSR (N.S.)* 73, (1950), 1117–1120
- [GKM] Gesztesy, F., Kiselev A.; Makarov, K. Uniqueness results for matrix-valued Schrodinger, Jacobi, and Dirac-type operators. *Math. Nachr.* 239/240 (2002), 103–145.
- [Ka] Kato, T. *Perturbation theory for linear operators*. Springer-Verlag, Berlin, 1995
- [KK1] Kargaev, P.; Korotyaev, E. Inverse Problem for the Hill Operator, the Direct Approach. *Invent. Math.*, 129(1997), no. 3, 567-593
- [KK2] Kargaev P.; Korotyaev, E. Effective masses and conformal mappings. *Comm. Math. Phys.* 169 (1995), no. 3, 597–625
- [K1] Korotyaev, E. Marchenko-Ostrovski mapping for periodic Zakharov-Shabat systems, *J. Differential Equations*, 175(2001), no. 2, 244–274
- [K2] Korotyaev, E. Inverse Problem and Estimates for Periodic Zakharov-Shabat systems, *J. Reine Angew. Math.* 583(2005), 87-115
- [K3] Korotyaev, E. Metric properties of conformal mappings on the complex plane with parallel slits, *Internat. Math. Res. Notices*, 10(1996), 493–503
- [K4] Korotyaev, E. Conformal spectral theory for the monodromy matrix, preprint 2006
- [K5] Korotyaev, E. Inverse resonance scattering on the real line, *Inverse Problems*, 21(2005), 1-17
- [Kr] Krein, M. The basic propositions of the theory of λ -zones of stability of a canonical system of linear differential equations with periodic coefficients. In memory of A. A. Andronov, pp. 413–498. *Izdat. Akad. Nauk SSSR, Moscow*, 1955
- [MO] Marchenko V., Ostrovski I. A characterization of the spectrum of the Hill operator. *Math. USSR Sb.* 26, 493-554 (1975)
- [Mi] Misura, T. Properties of the spectra of periodic and anti-periodic boundary value problems generated by Dirac operators. I,II, *Theor. Funktsii Funktsional. Anal. i Prilozhen*, (Russian), 30 (1978), 90-101; 31 (1979), 102-109
- [Po] Potapov, V. The multiplicative structure of J -contractive matrix functions. (Russian) *Trudy Moskov. Mat. Obšč.* 4, (1955). 125–236
- [RS] M. Reed ; B. Simon. *Methods of modern mathematical physics. IV. Analysis of operators*. Academic Press, New York-London, 1978
- [YS] Yakubovich, V., Starzhinskii, V.: *Linear differential equations with periodic coefficients*. 1, 2. Halsted Press [John Wiley & Sons] New York-Toronto, 1975. Vol. 1, Vol. 2

[Z] Zworski M., : Distribution of poles for scattering on the real line, J. Funct. Anal. 73, 277-296, 1987